



24 (C) REGISTRANT:
 Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

Alion™ Herbicide

Application in citrus groves to trees established less than one year

EPA REG. No. 264-1106

SLN No. FL-110004

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN FLORIDA

DIRECTIONS FOR USE

It is a violation of State and Federal Law to use this product in a manner inconsistent with its labeling. Persons using this product must comply with all applicable directions, restrictions, and precautions found on this labeling and that of the label of the federally registered product upon which this amendment is based. This labeling must be in the possession of the user at the time of pesticide application.

ALION™ Herbicide

Active Ingredient:

Indaziflam19.05%

Other Ingredients:80.95%

APPLICATION DIRECTIONS FOR USE IN RECENT PLANTED CITRUS GROVES

Only apply Alion Herbicide in citrus groves where the soil is completely settled around citrus trees and there are not open channels or depressions in the soil that would allow the product to move into the root zone through open channels. Alion Herbicide may be used around trees have been established for a minimum of one year after transplanting or in trees planted a minimum of one month provided the following conditions exists:

- The transplanted trees were potted plants (such as citripots) and not bare-rooted
- The trunks are protected from spray contact by nonporous wraps, grow tubes or waxed containers.
- The trees are actively growing and exhibiting good health and vigor.
- Application is made with broadcast equipment delivering a uniform spray pattern. Spot spraying is not allowed.

Avoid direct or indirect spray contact with crop foliage, green bark, roots or fruit as it may cause localized crop injury or death. Only the trunks of trees transplanted more than one year may be sprayed with Alion Herbicide if the trunk is callused, mature brown bark. Contact of Alion Herbicide with tissues other than mature brown bark can result in serious damage or plant death.

Citrus trees may be transplanted into soil previously treated with Alion Herbicide 1 month or more after the last application provided potted trees (such as citripots) are used.

Citrus Crops: calamondin, citron, citrus hybrids, grapefruit, kumquat, lemon, lime, orange, pummelo, tangelo, tangerine (mandarin) and satsuma mandarin

Dose Rate Chart for Citrus Groves

	Alion Herbicide (fl oz/A Broadcast)
Any soil texture	5.0 – 6.5 fl oz of product/treated A (0.065 to 0.085 lb ai/A) ¹

Notes and Restrictions:

¹ Choose the higher rate for longer weed control or situations with heavy weed debris.

Do not apply more than a total of 10.3 fl oz of product/treated A (0.134 lb ai/A) per year.

When making more than one application per year, allow a minimum of 90 days between applications.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937) or
 Visit our worldwide web site at www.bayercropscienceus.com
 As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).