



Supplemental Label

MANZATE PRO-STICK FUNGICIDE

EPA Reg. No. 70506-234

For Use on Broccoli, Cabbage, Lettuce (Head and Leaf), and Peppers

THIS SUPPLEMENTAL LABEL EXPIRES ON December 31, 2017 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

Read the label affixed to the container for MANZATE PRO-STICK, EPA Reg. No. 70506-234 before applying. Carefully follow all precautionary statements and applicable use directions.

This Supplemental Label contains new or supplemental instructions for use of this product which do not appear on the package label. Follow the instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the container labeling before buying or using MANZATE PRO-STICK. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of Limitation of Warranty and Liability on the container label.

USE RATES AND APPLICATION TIMINGS

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS/ACRE	TIMING/INTERVALS	RESTRICTIONS/COMMENTS
Broccoli	Alternaria Leaf Spot Downy Mildew	1.6 – 2.1 lb/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10-day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 12.8 lbs product (9.6 lbs mancozeb) /A per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days. Do not apply this product with a U-boom device. Aerial applications of Manzate

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS/ACRE	TIMING/INTERVALS	RESTRICTIONS/COMMENTS
				Pro-Stick on broccoli requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N, R, or P filter, NIOSH approval prefix TX-84-A.
Cabbage	Alternaria Leaf Spot Downy Mildew	1.6 – 2.1 lb/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10-day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 12.8 lbs product (9.6 lbs mancozeb)/A/season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days. Do not apply this product with a U-boom device. Aerial applications of Manzate Pro-Stick on cabbage requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N, R, or P filter, NIOSH approval prefix TX-84-A.
Lettuce (Head, Leaf)	Anthracnose Downy Mildew	1.6 – 2.1 lb/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10-day intervals, if needed. Use higher specified rates when conditions favor disease.	Remove residues from head lettuce by stripping and trimming. In California, do not apply more than 8.5 lbs product (6.4 lbs mancozeb)/A/ crop and do not apply within 14 days of harvest. In states other than California, do not apply more than 12.8 lbs product (9.6 lbs mancozeb)/A/crop and do not apply within 10 days of harvest. For all states: minimum retreatment interval is 7 days. Do not apply this product with a U-boom device.

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS/ACRE	TIMING/INTERVALS	RESTRICTIONS/COMMENTS
				Aerial applications of Manzate Pro-Stick on lettuce requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N, R, or P filter, NIOSH approval prefix TX-84-A.

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS/ACRE	TIMING/INTERVALS	RESTRICTIONS/COMMENTS
Pepper	Anthracnose Bacterial Spot (<i>Xanthomonas axonopodis</i> pv. <i>vesicatoria</i>) Cercospora Leaf Spot (Frogeye Spot) Phytophthora Blight Ripe Rot	West of the Mississippi: 1.6 – 2.1 lb/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10-day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 12.8 lbs product (9.6 lbs mancozeb)/A/season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days. Do not apply this product with a U-boom device. Aerial applications of Manzate Pro-Stick on pepper requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N, R, or P filter, NIOSH approval prefix TX-84-A.
		East of the Mississippi: 1.6 – 3.2 lb/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10-day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 19.2 lbs product (14.4 lbs mancozeb)/A/season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days. Do not apply this product with a U-boom device. Aerial applications of Manzate Pro-Stick on pepper requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N, R, or P filter, NIOSH approval prefix TX-84-A.

Rev08/15/2014