



FIFRA

Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
NEW YORK**

Caparol® 4L

**EPA Reg. No. 100-620
EPA SLN No. NY-140007**

For Control of Certain Weeds in Carrot and Cilantro

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2019

Active Ingredient:

Prometryn: 2,4-bis(isopropylamino)-6-(methylthio)-s-triazine 44.4%

Other Ingredients 55.6%

Total:.....100.0%

Caparol 4L contains 4 lbs. active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Caparol 4L label.

This label must be in the possession of the user at the time of pesticide application.

CARROT

Caparol 4L can be applied preemergence and or postemergence over the top to carrot. Read and follow all directions for use in carrot.

Use Precautions for Pre and Postemergence Applications in Carrot

Apply up to three applications of Caparol 4L at the rate of 2 - 4 pt/A per application.

Do not exceed one preemergence at up to 4 pt/A and two postemergence applications each at up to 2 pt/A or one postemergence application at up to 4 pt/A per crop cycle.

Do not exceed 8 pt/A of Caparol 4L per crop cycle

Make postemergence applications through the 6 leaf stage of carrot development

Do not apply within 30 days of harvest

Make uniform applications of the herbicide in a minimum of 20 gallons of water per acre.

When applying to emerged weeds add 2 qt of a nonionic surfactant (NIS) or wetting agent (approved for intended use) to 100 gal of spray mixture (0.5%) v/v or 1 gal of a non-phytotoxic crop oil concentrate (COC) containing 15-20% approved emulsifier to 100 gal of spray mixture (1% v/v).

Rotational Crops

The following crops may be seeded 5 months after applying no more than 4 pt Caparol 4L per acre on carrots: cabbage, carrots, celeriac, celery, Chinese celery, Florence fennel, corn, cotton, dill, fennel, okra and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

CILANTRO

Caparol 4L can be applied post planting, preemergence to cilantro. Read and follow all directions for use in cilantro.

Use Precautions for Preemergence Applications in Cilantro

Make one application of Caparol 4L at the rate of 2 – 3.2 pt/A.

Use the lower rates on coarse-textured soils and soils low in organic matter; use the higher rates on fine-textured soils and soils high in organic matter.

Do not exceed 3.2 pt/A of Caparol 4L per crop cycle.

Do not apply within 30 days of harvest.

Make uniform applications of the herbicide in a minimum of 20 gallons of water per acre.

Do not use on sand or loamy sand soil.

Rotational Crops

The following crops may be seeded 5 months after applying no more than 3.4 pt Caparol 4L per acre on cilantro: cabbage, carrots, celeriac, celery, Chinese celery, Florence fennel, cilantro, corn, cotton, dill, fennel, okra and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

Caparol® 4L is a trademark of Syngenta

Section 24(c) Registrant:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No: NY-140007

Label Code: NY0620071AA1014