



Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

ASSAIL® 30 SG INSECTICIDE FOR USE ON ALFALFA GROWN FOR SEED

This label is valid until December 31, 2019 or until otherwise amended, withdrawn, cancelled or suspended.

ACTIVE INGREDIENT: acetamiprid

(E)- N₁ -[(6-chloro-3-pyridyl)methyl]-N₂-cyano-N₁-methyl acetamidine.....30.0% by wt.

OTHER INGREDIENTS:70.0% by wt.

TOTAL100.0%

EPA Reg. No. 8033-36-70506

EPA SLN No. ID-090014

ENVIRONMENTAL HAZARDS:

This product is toxic to wildlife. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treated area. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not contaminate water used for irrigation or domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on this supplemental label and the main EPA registered label.

Spray volume: Apply in a minimum finished spray volume of 20 gallons per acre by ground and 10 gallons per acre by air. To achieve optimum pest control, it is important to obtain thorough and uniform spray coverage.

For use under this SLN label, do not apply this product through any type of irrigation system.

USE	RATE/ACRE	SPECIFIC DIRECTIONS
Lygus Bugs	2.5 – 4.0 oz (0.05 – 0.075 lb ai)	Begin application when treatment thresholds have been reached. Use the high rate under heavy insect pressure and for longer residual control. Do not apply ASSAIL 30 SG while bees are actively visiting the treated area.

- Do not make more than 3 applications per season.
- Do not apply more than once every 14 days.
- Do not exceed a total of 0.225 lbs. acetamiprid active ingredient (12.0 ozs of ASSAIL 30 SG product) per acre per growing season
- Do not apply less than 7 days before harvest (7 day PHI).



Special Local Need

SPECIAL CROP USE RESTRICTIONS:

Do not cut current year's treated alfalfa seed crop for hay or forage. Do not graze current year's treated alfalfa crop. Do not use treated alfalfa seed for sprouting.

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed. This restriction pertains to, but is not limited to, green chop, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or re-growth. All seed screenings must be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes.

Any seed from a field treated with this pesticide product must bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling must contain the following statement:

“This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration and other regulatory agencies.”

RESISTANCE MANAGEMENT:

Acetamiprid is the active ingredient in ASSAIL 30 SG Insecticide. It is a member of a class of chemicals known as neonicotinoids and within the mode of action Group 4A. Rotating ASSAIL 30 SG Insecticide with insecticides of a different mode of action (other than Group 4A insecticides) may delay or prevent development of resistance and cross-resistance to ASSAIL and other Group 4A insecticides. Avoid making more than two (2) consecutive applications of ASSAIL 30 SG Insecticide before rotating to an alternative mode of action insecticide. Foliar applications of ASSAIL 30 SG Insecticide should be avoided on crops treated with a Group 4A seed treatment or soil-applied insecticide until a foliar application of a non-Group 4A insecticide (insecticide with a different mode of action) has been applied between these applications. The use of ASSAIL 30 SG Insecticide should conform to the resistance management guidelines established in your area. Consult your agricultural advisor, PCA, university or extension personnel for recommended pest and resistance management practices for your area. Use recommended IPM practices in your pest management system. Use of rates below the minimum rate listed for each particular insect pest may enhance the development of resistance and should be avoided.

Assail® is a registered trademark of Nippon Soda Co., Ltd.

(Rev. 3/18/14)