



# Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN MAINE,  
MARYLAND, MICHIGAN AND NEW JERSEY

## Assail<sup>®</sup> 30SG Insecticide EPA Reg. No. 8033-36-70506

### Additional Use Directions for Blueberries

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Pest	Dosage per acre	Specific Directions
Spotted Winged Drosophila (SWD) Larvae	5.3 oz. (0.1 lbs. ai)	Thorough coverage is important to obtain optimum control. Begin application when adults are present and fruit is ripening or already ripe. It is recommended to tank mix with an adulticide.

#### Restrictions and Precautions:

- Do not make more than 5 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest (PHI = 1 day).
- Do not exceed a total of 0.5 lbs. active ingredient (26.7 oz.) per acre per growing season.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

Rev. 03/20/2014