

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ MATRIX® SG HERBICIDE

EPA Reg. No. 352-768

FOR PREMERGENCE CONTROL OF ADDITIONAL GRASS AND BROADLEAF WEEDS IN LABELED TREE AND VINE CROPS IN THE STATES OF ARIZONA, CALIFORNIA, OREGON AND WASHINGTON

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such non-labeled use.

DuPont™ MATRIX® SG Herbicide is recommended for the preemergence control of annual sowthistle, annual bluegrass, whitestem filaree, London rocket, panicle willowweed, prickly lettuce, yellow sweetclover, white clover, burclover, common chickweed, redmaids, coast fiddleneck, Italian ryegrass, swinecress and malva/cheeseweed in labeled tree and vine crops.

Apply MATRIX® SG Herbicide at the rate of 4.0 ounces product per acre prior to weed emergence.

PRECAUTIONS:

Prior to using MATRIX® SG Herbicide for preemergence control of the weeds on this label, consult the MATRIX® SG Herbicide section 3 federal label and read instructions under the heading of "citrus fruit, stone fruit, tree nut, pome fruit and

grapes." It is important to use a postemergence/contact herbicide such as glyphosate or other registered postemergence herbicides to control any emerged weeds in order to obtain the best residual weed control from MATRIX® SG Herbicide. Also, depending on your weed complex, you may need to include another residual herbicide as a tankmix treatment with MATRIX® SG Herbicide.

Summer annual weed species may not be controlled or suppressed by MATRIX® SG Herbicide, depending on the time of germination throughout the summer months.

Important - Before using DuPont Matrix ®SG Herbicide read and follow all applicable directions, restrictions and precautions on the EPA-registered label.

RESTRICTIONS:

Do not apply more than 4.0 oz MATRIX® SG Herbicide per year on a broadcast acre basis. When applied as a banded treatment (50% band or less), MATRIX® SG Herbicide may be applied twice per year.

Allow a minimum of 30 days between applications

The field re-entry interval (REI) is 4 hours.

Preharvest interval (PHI): Last application (days to harvest) – Citrus Fruit 3 days, Pome Fruit 7 days, Tree Nuts 14 days, Stone Fruit 14 days and Grapes 14 days.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805