



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709

ALIETTE® WDG Fungicide

For Aerial Application on Hops for Downy Mildew Control

EPA Reg. No. 264-516

EPA SLN. No. OR-040018

FIFRA 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Aliette® WDG Fungicide and this Section 24c label before applying. Do not use in manner inconsistent with applicable directions, restrictions, and precautions provided on the label affixed to the container for Aliette® WDG Fungicide and this Section 24c label. Use of Aliette® WDG Fungicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Aliette® WDG Fungicide.

THIS LABEL IS VALID UNTIL DECEMBER 31, 2017 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

ALIETTE® WDG Fungicide

ACTIVE INGREDIENT: Aluminum tris (O-ethyl phosphonate).....	80.0%
INERT INGREDIENTS:	20.0%
TOTAL:	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye injury. Avoid contact with skin, eyes or clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.

SPECIFIC DIRECTIONS FOR USE

Crop/Site:	Hops
Target Pest:	Downy Mildew (<i>Pseudoperonospora humili</i>)
Rate:	2.5 lbs / Acre.
Dilution Rate:	Use sufficient water to insure complete coverage of foliage. Do not use less than 10 GPA for aerial application.
Method of Application:	Foliar spray
When to Apply:	Apply ALIETTE® WDG Fungicide as a foliar spray. When conditions are favorable for disease development (warm and humid), applications should be made as follows: (1) when shoots are 6-12 inches high; (2) after training when vines are 5-6 feet tall; (3) approximately three weeks after the second application; and (4) during bloom. Aerial application may result in reduced control due to lack of canopy penetration and coverage. It should only be used in situations where ground application is impossible.
Restrictions and Limitations:	Do not apply within 24 days of harvest (PHI = 24 Days). Do not apply more than 10 lbs/Acre per growing season. Do not feed hop refuse to animals.

Note: Bayer CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE WDG Fungicide or if unbuffered ALIETTE WDG Fungicide is applied to foliage with copper residues. Do not tank-mix with copper fungicides.

When applied prior to or after copper fungicides, the pH of ALIETTE WDG Fungicide should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE WDG Fungicide) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE WDG Fungicide).

Adjuvants which enhance pesticide penetration may cause phytotoxicity when mixed with ALIETTE® WDG.

Mixing ALIETTE WDG Fungicide with surfactants or foliar fertilizers is not recommended. When ALIETTE WDG Fungicide is used in combination with any other pesticide, conduct a crop sensitivity test by spraying a few plants. Evaluate the sprayed plants 3-7 days after application for adverse effects.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).