



## Section 2(ee) Recommendation

Date: March 18, 2015  
Product: Velista™, EPA Reg. No. 100-1534  
Use: Addition of Diseases as Pests Controlled on Turfgrasses  
State: OR

### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

| Disease   | Application Site | Rate oz/1,000 sq ft | Application Interval (Days) | Comments  |
|---|------------------|---------------------|-----------------------------|---|
| Rapid Blight ( <i>Labyrinthula terrestris</i> ) | All Turfgrass    | 0.5                 | 14                          | Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz product/1,000 sq ft/year.  |
| Fairy Ring ( <i>Basidiomycetes</i> )            | All Turfgrass    | 0.5-0.7             | 14-28                       | Apply as part of a preventative program. Rotate with Heritage® and Briskway® fungicides. Do not apply more than 2.2 oz product/1,000 sq ft/year.                                      |
| Leaf and Sheat Spot ( <i>Rhizoctonia zeae</i> ) | All Turfgrass    | 0.3-0.5             | 14                          | Use as part of a preventative program in tank mixture and/or rotation with Daconil Action™, Heritage and Briskway fungicides. Do not apply more than 2.2 oz product/1,000 sq ft/year. |
| Spring Dead Spot ( <i>Ophiosphaerella</i> spp.) | All Turfgrass    | 0.7                 | 28                          | Make two applications on a 28 day interval prior to winter dormancy. Do not apply more than 2.2 oz of product/1,000 sq ft/year.   |

| <b>Disease</b>  | <b>Application Site</b> | <b>Rate oz/1,000 sq ft</b> | <b>Application Interval (Days)</b> | <b>Comments</b>  |
|---|-------------------------|----------------------------|------------------------------------|--|
| SummerPatch<br>( <i>Magnaporthe poae</i> )                              | All Turfgrasses         | 0.3-0.5                    | 14-28                              | During periods of high disease pressure, tank mix with a fungicide that is registered for the control of summer patch. Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz of product/1,000 sq ft/year. |
| Brown Ring Patch<br>( <i>Waitea circinata</i><br><i>var circinata</i> ) | All Turfgrasses         | 0.5                        | 14                                 | Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz of product/1,000 sq ft/year.  |

Velista™, Briskway®, Daconil Action®, Heritage® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2015 Syngenta

2(ee) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, NC 27419-8300

Expiration Date: December 31, 2019

Label Code: OR1534179AC0315