



# Supplemental Label

## Manzate® Pro-Stick™ Fungicide

EPA Reg. No. 70506-234

### Directions for Use on Ferns (not for this use in California)

#### DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its label. This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DISEASES	RATE OF MANZATE PRO- STICK	DIRECTIONS
Botrytis Cercospora Curvularia Cylindrosporium Glomerella Phyllosticta Taphrina	1 – 2 lbs/A in 100 gallons water	<p>Apply as a thorough coverage spray. Begin application at first sign of disease and repeat at 7 to 10 day intervals or as needed; use shorter interval during periods of frequent rains or when severe disease conditions persist.</p> <p>Aerial application: For aerial applications made to field-planted ornamentals, apply 1 to 2 lbs. per acre; a minimum rate of 5 gals of spray per acre should be used during aerial applications.</p> <p>Manzate Pro-Stick may be used alone or in combination with other fungicides as maintenance spray. Use higher rate and shorter intervals during periods of excessive wetness and rapid growth.</p> <p>Treated plants must not be used for food or feed purposes.</p>

**Note:** Plant sensitivities to Manzate Pro-Stick have been found to be acceptable in specific genera and species listed on this label, however, phototoxicity may occur. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test each one for sensitivity to Manzate Pro-Stick. Neither the manufacturer nor seller has determined whether or not Manzate Pro-Stick can be safely used on ornamental or nursery plants not listed on this label. The user should determine if Manzate Pro-Stick can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e. bedding plants, foliage, etc., and observe 7 to 10 days for symptoms of phytotoxicity prior to commercial use.