

- Date: February 17, 2004
- Products: Dual II MAGNUM[®] Herbicide (EPA Reg No. 100-818)
Dual II MAGNUM[®] SI Herbicide (EPA Reg No. 100-829)
Dual IIG MAGNUM[®] Herbicide (EPA Reg No. 100-910)
Dual MAGNUM[®] Herbicide (EPA Reg No. 100-816)
Bicep II MAGNUM[®] Herbicide (EPA Reg No. 100-817)
Bicep Lite II MAGNUM[®] Herbicide (EPA Reg No. 100-827)
Liberty[®] Herbicide (EPA Reg No. 264-660)
Callisto[®] (EPA Reg No. 100-1131)
- Use: Liberty at 20 oz/A may be applied post-emergence following a full labeled rate of Dual MAGNUM or Bicep MAGNUM brand herbicides applied at or before planting for improved control of Woolly Cupgrass in LibertyLink[®] corn. Callisto at 3 fl oz/Acre may be applied in tank mixture with Liberty for post-emergence broadleaf weed control.
- States: Illinois, Iowa, Minnesota, South Dakota, and Wisconsin

DIRECTIONS FOR USE

It is a violation of federal law to use these products in a manner inconsistent with their labeling.

For improved control of woolly cupgrass, apply Liberty at 20 oz/A post-emergence following a full labeled rate of a Dual MAGNUM or Bicep MAGNUM brand herbicide applied at or before planting. Liberty must be applied to woolly cupgrass prior to tillering and when cupgrass is less than 3 inches in height. Apply Liberty in 20-30 gallons of water per acre. Callisto at 3 fl oz/A per acre may be tank mixed with Liberty for improved postemergence broadleaf weed control. Always add spray grade AMS at 3 lbs./A to the spray solution when applying Liberty alone (see the Liberty label for additional adjuvant instructions). AMS at 3 lbs./A plus a non-ionic surfactant at 0.25% v/v (1 qt/100 gal of water) is recommended when Callisto is applied in a tank mix combination with Liberty. No other adjuvants may be used with Liberty and Callisto tank mix combinations.

Notes: 1) Liberty may only be applied to corn specifically designated as having the LibertyLink trait. 2) Liberty may be applied to LibertyLink corn up to 24 inches tall or in the V-7 stage of growth. 3) Do not apply Liberty within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder.

Refer to the Callisto and Liberty EPA approved container labels for complete directions for use and all applicable restrictions and precautions.

This labeling should be in the possession of the user at the time of pesticide application.

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by EPA.

Syngenta Crop Protection, Inc.
Greensboro, NC 27419

Dual II MAGNUM[®], Dual II MAGNUM[®] SI, Dual IIG MAGNUM[®], Dual MAGNUM[®], Bicep II MAGNUM[®], Bicep Lite II MAGNUM[®] and Callisto[®] are trademarks of a Syngenta Crop Protection, Inc..

Liberty and LibertyLink are registered trademarks of Bayer Cropscience.

IL1131022AC0204

ILCallistoLibertyLinkCornSec2ee 2-17-04