



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

GEM™ 500 SC Fungicide

For Use in Artichoke to control Ramularia leaf spot

EPA Reg. No. 264-826

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

ARTICHOKE (GLOBE)			
Disease Controlled	Rate fl. oz./Acre	Application Timing	Notes
Ramularia leaf spot <i>(Ramularia spp.)</i>	3.0 – 3.8	Begin applications when conditions are favorable for diseases but before infection.	Use the higher rates when disease pressure is severe. Apply on a 7 to 10-day spray schedule.
Restrictions			
Do not apply more than 7.6 fl. oz. of Gem 500 SC Fungicide per acre per season. Gem 500 SC Fungicide maybe applied up to the day of harvest (0-day preharvest interval). To limit the potential for development of disease resistance alternate each application of Gem 500 SC Fungicide with a non-Group 11 containing fungicide.			

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information, Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*