



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Poncho®/VOTiVO®

Add Sugarcane Aphid (*Melanaphis sacchari*)
for Commercial Sorghum Seed Treatment.

EPA Reg. No. 264-1109

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, IOWA, KANSAS, LOUISIANA, MISSOURI, MISSISSIPPI, NEBRASKA, NEW MEXICO, OKLAHOMA, TENNESSEE AND TEXAS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

COMMERCIAL SORGHUM SEED TREATMENT

Pest	Rate of Poncho/VOTiVO	Remarks
Sugarcane Aphid	Apply 6.13 fl oz per 100 lbs seed	Apply prior to planting as a slurry treatment to provide early season protection of seedlings. Ensure thorough coverage.

RESTRICTIONS:

Regardless of type of application (seed treatment or foliar), do not apply more than 0.02lbs (sorghum) of active ingredient clothianidin per acre per season.

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with Poncho/VOTiVO must be labeled with the following statements:

- This seed has been treated with PONCHO/VOTiVO, which contains **40.3%** clothianidin **and 8.1%** *Bacillus firmus* I-1582.
- Do not use treated seed for food, feed, oil production or ethanol production.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with Poncho/VOTiVO

- Regardless of type of application (seed treatment or foliar), do not apply more than 0.02lbs (sorghum) of active ingredient clothianidin per acre per season.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- This compound may be highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals.

- Cover or collect treated seeds spilled during loading.
- For Cotton, Sweet corn, and Sorghum: **Immediate Plant-back:** Cereal grains, Field corn, popcorn, sweet corn, cotton, rapeseed, soybeans, canola, sorghum, root and tuber vegetables, **30-Day Plant-back:** Grasses, non-grass animal feeds, dried beans, **4- Month Plant-back:** Any crop without an earlier plant-back interval

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*