



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
http://www.bayercropscienceus.com

SCALA™ brand SC Fungicide

EPA Reg. No. 264-788

For use on Almond, Pistachio, Bulb Vegetables, Grapes, Pome Fruit, Stone Fruit, Strawberry

Supplemental Label

SCALA™ brand SC Fungicide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for SCALA™ brand SC Fungicide.

ALMONDS, PISTACHIOS

Table with 4 columns: CROP, DISEASE, USE RATE (FL. OZ. PER ACRE, LBS. A.I. PER ACRE), and USE INFORMATION. Rows include Almond/Pistachio with diseases like Brown rot blossom blight, Jacket rot, Green fruit rot, Shot hole, Alternaria leaf spot, Botryosphaeria blight, Anthracnose, Scab, and Blossom & shoot blight.

BULB VEGETABLES

Table with 4 columns: CROP, DISEASE, USE RATE (FL. OZ. PER ACRE, LBS. A.I. PER ACRE), and USE INFORMATION. Rows include Onion, Green, Dry Bulb, Welsh, Garlic, Great-Headed, Leek, Shallot with diseases like Botrytis leaf blight, Neck rot, and Purple blotch.

GRAPES

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Grape	<b>Gray mold</b> ( <i>Botrytis cinerea</i> )	<b>SCALA™ SC Alone</b>		Use lower rate only in a tank mix with another fungicide active against <i>Botrytis</i> . Applications should be made at the critical timings for <i>Botrytis</i> control. Typically, first applications are made at early bloom, and/or berry touch to bunch closure, veraison, and pre-harvest. Use sufficient water to ensure penetration of the canopy and coverage of the flowers or bunches.  Do not apply more than 36 fl. oz./A (1.4 lbs. a.i./A) of SCALA™ SC per crop. Do not apply within 7 days of harvest.
		18	0.70	
		<b>SCALA™ SC In tank mixes</b>		
		9 - 18	0.35 - 0.70	

POME FRUITS

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Apple Pear (All varieties) Pear, Oriental Crabapple Loquat Mayhaw Quince	<b>Scab</b> ( <i>Venturia</i> spp.)	<b>SCALA™ SC Alone</b>		Under conditions favorable for disease development, shorten the spray intervals and/or use the high rate. Use lowest rate only in a tank mix with another fungicide active against scab. For optimum results, begin applications at green tip or as soon as crop and/or environmental conditions become favorable for disease development. Applications should be made on 7 day or longer intervals depending upon disease conditions. Do not apply more than 40 fl. oz./A (1.6 lbs. a.i./A) of SCALA™ SC per crop. Do not apply within 72 days of harvest.
		7 - 10	0.27 - 0.39	
		<b>SCALA™ SC In tank mixes</b>		
		5 - 10	0.19 - 0.39	

STONE FRUITS  
(Except Cherries)

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Apricot Nectarine Peach Plum (All varieties) Plum, Chickasaw Plum, Damson Plum, Japanese Plum, Stanley Plumcot Prune	<b>Brown rot blossom blight</b> ( <i>Monilinia</i> spp.)	<b>SCALA™ SC Alone</b>		For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Apply when bud tissue is susceptible to infection (i.e., pink, white or red bud). If conditions favorable for disease development persist or recur, apply at full bloom or at petal fall. Pre-harvest applications should be made on 7 day intervals or longer. Do not apply more than three applications of SCALA™ SC alone or more than 54 fl. oz./A (2.1 lbs. a.i./A) of SCALA™ SC per crop. Do not apply within 2 days of fruit harvest or make more than 2 applications of a Group 9 fungicide within 30 days of harvest. <b>Do not use on cherries.</b>  For shot hole, scab, and brown fruit rot control, use the lower rate only in a tank mix with a fungicide registered for use against these diseases.
		9 - 18	0.35 – 0.70	
	<b>SCALA™ SC Alone</b>			
	18	0.70		
	<b>SCALA™ SC In Tank Mixes</b>			
	9 - 18	0.35 - 0.70		
	<b>Jacket rot, Green fruit rot</b> ( <i>Botrytis cinerea</i> )			
	<b>Shot hole</b> ( <i>Wilsonomyces carpophilus</i> )			
	<b>Scab</b> ( <i>Cladosporium</i> spp.)			
	<b>Brown fruit rot</b> ( <i>Monilinia fructicola</i> )			

STRAWBERRIES

CROP	DISEASE	USE RATE		USE INFORMATION
		FL. OZ. PER ACRE	LBS. A.I. PER ACRE	
Strawberry	<b>Gray mold</b> ( <i>Botrytis cinerea</i> )	<b>SCALA™ SC Alone</b>		Use lower rate only in a tank mix with another fungicide active against gray mold.  For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications from pre-bloom to harvest on a 7 to 14 day interval depending upon disease conditions.  Do not apply more than 54 fl. oz./A (2.1 lbs. a.i./A) of SCALA™ SC per crop. Always follow resistance management directions listed on this label and the recommendations for your area. Do not apply within 1 day of harvest.
		18	0.70	
		<b>SCALA™ SC In tank mixes</b>		
		9 - 18	0.35 - 0.70	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

SCALA is a trademark of Bayer.