

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

For Agricultural or Commercial Use Only

CROP	PEST	RATE OF APPLICATION
Fruiting Vegetables (except Cucurbits) Leafy Vegetables Head, Stem and Leafy Brassica Vegetables Legume Vegetables	Symphylans	4.3 ounces per acre

DIRECTIONS FOR USE

This recommendation is to add a pest to the label for which control data is now available.

Apply Mustang Insecticide before, during, or after planting for protection from symphylans. Placement of Mustang Insecticide is critical to get plant protection and will depend on seed type and placement, or transplant placement. Successful protection from symphylans is obtained by having Mustang Insecticide-treated soil around or in contact with developing plant roots. Whatever application method is used must result in proper placement of Mustang Insecticide to get plant protection. Successful application methods include (but may not be limited to) the following:

Direct-seeded crops: Mustang Insecticide should be applied in a narrow band (3-6 inches) and incorporated throughout the seed line. Incorporation should result in treated soil around and below (1-3 inches) the seed. Mechanical and sprinkler incorporation have been successful, but mechanical incorporation has been more consistent. Where more than one seed line per bed are planted, multiple bands or bed top applications will be necessary. Use as narrow a band as practical.

Transplanted crops: Apply Mustang Insecticide to the transplant row prior to transplanting in as narrow a band as practical (but not less than 4-6 inches) and mechanically incorporate to a depth of 4-6 inches. Mustang Insecticide may also be applied as a soil drench at the time of transplanting. Mustang insecticide has also given plant protection from symphylans when applied to the transplant row with two shanks four inches apart and six inches deep. Mustang Insecticide may also be applied with a spray blade so a barrier of treated soil immediately below the transplant is obtained.

RESTRICTIONS

Follow phi (pre-harvest interval) restrictions as given on the Mustang Insecticide label.

A maximum of 0.3 lb active ingredient may be applied per acre per season. Do not make applications of Mustang Insecticide less than 7 days apart (except for Legume Vegetables which can be made 5 days apart).

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG INSECTICIDE LABEL (EPA Registration No. 279-3126) MUST BE FOLLOWED.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.