

**SUPPLEMENTAL LABELING  
FOR DISTRIBUTION AND USE ONLY WITHIN MARYLAND**



# Acramite® -50WS

**MITICIDE IN WATER SOLUBLE BAGS  
EPA REG. NO. 400-503      EPS SLN. NO. MD-050002**

Active Ingredients: (% by weight)  
Bifenazate: Hydrazine carboxylic acid, 2-(4-methoxy-  
[1,1'-biphenyl]-3-yl) 1-methylethyl ester ..... 50%  
Inert Ingredients ..... 50%  
Total..... 100%

## KEEP OUT OF REACH OF CHILDREN CAUTION

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AERIAL APPLICATION TO CUCURBIT AND FRUITING VEGETABLES TO CONTROL TWO-SPOTTED SPIDER MITES::

#### USE INSTRUCTIONS

CROP	AMOUNT ACRAMITE-50WS PER ACRE	MINIMUM GALLONS PER ACRE BY AIR	TOTAL NUMBER OF SPRAYS PER YEAR	REI
Cucurbit Vegetables* Fruiting Vegetables**	1.0 lb.	5	1	12 hours

\*Cucumber; edible gourd (includes hechima and Chinese okra); muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honey dew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon); pumpkin; squash (includes crookneck, scallop squash, straightneck squash, vegetable marrow, zucchini); winter types (includes butternut squash, calabaza, hubbard squash; acorn squash and spaghetti squash); watermelon

\*\*Eggplant; groundcherry; pepino, tomatillo; peppers (includes bell pepper, chili peppers, cooking peppers, pimento and sweet pepper); tomatoes.

**NOTE: Observe 3 day pre-harvest interval (PHI) for all crops listed.**

**DRIFT:** Employ all spray control measures to limit drift from the application site that may contact non-target sites such as non-target crops, woodlands, aquatic areas, and wetland areas. Avoiding spray drift is the responsibility of the applicator. The applicator must use all available measures to control drift. Do not spray if wind speeds are >10 mph. Use caution when conditions are favorable for drift such as high temperatures, drought, and low relative humidity. Do not apply when a temperature inversion exists.

Only registered crops may be rotated in a treated field within 30 days of the final application to avoid uptake of residues.

## THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER

**Follow all applicable directions, restrictions, and precautions on the EPA registered label.**

**This label must be in the possession of the user at the time of the pesticide application.**

**004**

ACRAMITE® is a Registered Trademark of Chemtura Corporation  
24c (SLN) REGISTRANT:  
Chemtura Corporation  
199 Benson Road  
Middlebury, CT 06749

EPA SLN NO. MD-050002