

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

GoalTender™

EPA Reg. No. 62719-447

For Control of Weeds in Non-Cropland

(For Distribution and Use Only Within the State of California)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for GoalTender™ herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of GoalTender according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for GoalTender.

Directions for Use

NON-CROP USE

Non-food producing, non-cultivated agricultural or non-agricultural areas, such as highway and utility rights-of-way, industrial sites, tank farms, storage areas, airports, fencerows, and farmsteads

General Information

GoalTender is recommended for postemergence and preemergence control of listed broadleaf weeds in non-crop areas.

Weeds Controlled Postemergence (weeds up to 4 inches high): Apply 1 to 4 pints (0.5 to 2 lb active) of GoalTender per broadcast acre. The lower rate in the rate range is recommended for control of susceptible weeds in the early postemergence stage, less than 4 inches in height. The higher rate (2 lb active) should be used for weeds up to 12 inches in height. Applications to weeds beyond the 4-inch stage may result in partial control.

cheeseweed (malva)
fiddleneck, coast
filaree, broadleaf
filaree, redstem
groundsel, common
henbit
minerslettuce
nettle, burning
pigweed, redroot
purslane, common
redmaids
shepherdspurse
sowthistle, annual

Weeds Controlled Preemergence: Apply 2.5 to 4 pints (1.25 to 2 lb active) per broadcast acre.

burclover
cheeseweed (malva)
fiddleneck, coast
filaree, broadleaf
filaree, redstem
groundsel, common
henbit
knotweed, prostrate
lambquarters, common
lettuce, prickly
pigweed, redroot
purslane, common
redmaids
rocket, london
shepherdspurse
sowthistle, annual

Timing and Method of Application

GoalTender should be applied in a minimum of 40 gallons of water per acre. Best preemergence results are achieved when spray is applied to a relatively weed free soil surface. The volume of water used should be increased as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

Tank Mixes With GoalTender

Important: Read and observe all label directions before using. When tank mixing, read and carefully follow all applicable use directions, precautions, and limitations on the respective product labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

Dosage

For preemergence control of susceptible grass and broadleaf weeds, a tank mixture of GoalTender with diuron (Karmex) or simazine can be applied. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

For postemergence control of susceptible grass and broadleaf weeds, a tank mixture with paraquat (Gramoxone) or glyphosate (Glyphomax) with GoalTender can be used. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

Specific Use Restrictions

In addition to the following, also observe General Use Restrictions listed on the label affixed to the container for GoalTender.

- Do not feed or allow animals to graze on any areas treated with GoalTender.

TMTrademark of Dow AgroSciences LLC

R204-026

EPA Accepted: 02/16/05

Initial printing.