

# Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger<sup>®</sup>

EPA Reg. No. 62719-73

### 2(ee) Recommendation<sup>†</sup>

For Distribution and Use Only in the State of Washington

### Use Recommendations for Exported Grass Hay

#### ATTENTION

<sup>†</sup> This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to container for Stinger<sup>®</sup> herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of Stinger is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

#### Directions for Use

This product bulletin should be followed if the target grass crop will be exported to clopyralid-sensitive destinations.

Timing of Application	Spring Cutting	Fall Cutting	Spring Cutting Subsequent Year <sup>1</sup>	Fall Cutting Subsequent Year <sup>1</sup>
before spring cutting	do not export	may be exported	may be exported	may be exported
after spring cutting	N/A	do not export	may be exported	may be exported
after fall cutting	N/A	N/A	may be exported	may be exported

N/A – not applicable

<sup>1</sup>If no clopyralid applications are made in subsequent year.

- Fall applications should be made while grass and weeds are actively growing.
- Fall applications should be made as close to last cutting as possible in order to reduce clopyralid residues in hay the following year.
- Adequate soil moisture, particularly with fall applications, will help weed control as well as reduce clopyralid residues in hay.
- In areas where three cuttings can be made, avoid exporting the first cutting after a clopyralid application.

<sup>®</sup>Trademark of Dow AgroSciences LLC

R043-031

Issued: 01/27/06

Initial printing