

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Control of Powdery Mildew on Artichoke

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Rally 40WSP (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Erysiphe cichoracearum</i>)	4	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage. Use a minimum of 30 gallons of spray volume per acre. Control may be reduced at low spray volumes or if spray coverage is not adequate. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 4 oz (0.25 lb) of Rally 40WSP (0.1 lb ai) per acre per application or 24 oz (1.5 lb) of Rally 40WSP (0.6 lb ai) per acre per season. • Do not make more than 6 applications per season. • Pre-harvest Interval: Do not harvest within 3 days of application • Re-treatment Interval: 14 days • Rotational Crop Restriction: Crops with registered uses may be planted immediately after the last treatment. Do not plant other crops within 30 days after the last application.

Resistance Management: Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R182-027

EPA accepted 09/11/09

Replaces R182-004