

## RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

# FIFRA 24(c) Special Local Need Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Lorsban<sup>®</sup> Advanced

EPA Reg. No. 62719-591

EPA 24(c) Special Local Need Registration SLN CA-080010  
For Distribution and Use Only in the State of California

### Control of Argentine Ants Infesting Grapes

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban<sup>®</sup> Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

#### Directions for Use

Apply 1.5 to 2 pints of Lorsban Advanced per acre in a minimum of 2.5 gallons of water per acre using a power-operated ground or hand-held sprayer. Maintain constant agitation of the spray tank. To obtain thorough coverage, spray the base of each vine, the vine stakes, and the surrounding soil to about 1 foot from the base of the vine. If a hand-held sprayer is used, 2 to 5 fl oz per vine is required. A non-petroleum-based spreader/sticker spray adjuvant may be used if previous experience indicates that it is safe under the intended conditions of use. Follow use directions for the spreader/sticker product.

#### Specific Use Precautions:

- Do not allow spray to contact fruit or foliage.
- Weed growth in the vine rows should be controlled to prevent ants from trailing to the vine cordons via weeds and to allow better spray coverage.
- Consult your extension service specialist or certified pest control advisor for use information specific to your area.

#### Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 76 days before harvest.
- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Do not make more than 3 applications per year.
- Maximum single application rate is 0.94 lb ai chlorpyrifos per acre

- For residue purposes, growers may treat grapes using Lorsban Advanced under the requirements of this SLN or Lorsban Advanced under the requirements listed in SLN CA-080009. Growers may not treat the same grape crop under the requirements of both SLNs, nor may they treat the same grape crop under the requirements of any SLN for Lorsban-4E.

**Expiration Date: December 31, 2019**

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R368-105

Approved: 10/13/14

Replaces R368-050