

# Supplemental Labeling



**Dow AgroSciences LLC      9330 Zionsville Road      Indianapolis, IN 46268-1054 USA**

**RESTRICTED USE PESTICIDE**  
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## Lorsban<sup>®</sup> Advanced

EPA Reg. No. 62719-591

**EPA 24(c) Special Local Need Registration SLN VA-090001  
For Distribution and Use Only in the State of Virginia**

### Insect Control Infesting Tobacco

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban<sup>®</sup> Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

### Directions for Use

Refer to product label for Lorsban Advanced for Use Precautions and Restrictions, Mixing and Application instructions.

Apply as a preplant broadcast spray to reduce the feeding damage caused by listed pests. Apply 24 to 48 hours before bedding and transplanting using a spray volume of 10 gallons per acre (gpa) or more. Incorporate immediately after application to a depth of 2 to 4 inches using suitable incorporation equipment.

Before broadcast application of Lorsban Advanced onto existing beds, knock down beds to final shape for transplanting. Use of PTO-driven implements that will incorporate Lorsban Advanced to a depth of 4 inches is recommended.

<b>Target Pests</b>	<b>Lorsban Advanced (quart/acre)</b>
cutworms flea beetles mole crickets root maggots wireworms	2

To control the above listed pests and suppress populations of rootknot nematodes in all tobacco growing regions, use Lorsban Advanced in a tank mix with Nemacur 3 at the rate of 2 quarts of Lorsban Advanced

plus 4 quarts of Namacur 3 nematicide per acre. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for Namacur 3 used in combination with Lorsban Advanced.

Apply the specified rate(s) to the soil surface in a spray volume of 10 gpa or more 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil to a depth of at least 4 inches using suitable equipment. Where the nematode species *Meloidogyne arenaria* or *M. javanica* are present or high populations of *M. incognita*, apply Telone® II soil fumigant at the recommended label rate.

**Specific Use Precautions and Restrictions:**

- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Do not make more than one application of Lorsban Advanced or other product containing chlorpyrifos per season.
- Maximum single application rate is 1.88 lb ai chlorpyrifos per acre.
- **Chemigation:** Do not apply through any type of irrigation equipment.

®Trademark of Dow AgroSciences LLC

R368-047

Approved: 04/01/09

Initial printing.