

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Vista[®] XRT

EPA Reg. No. 62719-586

For Control of Broadleaf Weeds including Prickly Pear Cactus in Rangeland and Permanent Grass Pastures

(For Distribution and Use in the State of Florida)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Vista[®] XRT herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Vista XRT according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

General: Broadcast apply Vista XRT[®] herbicide as a single treatment or as sequential postemergence treatment using ground or aerial application equipment. Vista XRT may be applied in tank mix combination with other foliar-applied herbicides labeled for use on rangeland and permanent pastures to control additional weeds and woody plants. Read and follow applicable use directions, precautions and limitations on each product label. Refer to the General Information section of this label for application instructions, including application timing, effect of temperature on herbicidal activity, application rate ranges, spray coverage, instructions for spot application and tank mixing.

Application Timing and Weeds Controlled or Suppressed: Apply as a broadcast treatment when weeds are actively growing, but prior to bud stage of weed growth. Use a total spray volume of 5 or more gallons per acre for ground broadcast or 3 or more gallons per acre by air. Only weeds emerged at the time of application will be controlled.

Tank Mixtures: For control of additional weeds and woody plants, Vista XRT may be tank-mixed with Remedy[®] herbicide at the indicated application rates.

Products in Tank Mix	Application Rates	Additional Weeds/Brush Controlled	
Vista XRT Remedy	4 fl oz/acre + 9 fl oz/acre	buttercup, hairy croton dogbane, hemp kochia lespedeza, sericea	marshelder ragweeds sunflower thistle, musk vetch
Vista XRT Remedy	6 fl oz/acre + 3/4 pt/acre	dandelion dock, curly dogfennel goldenrod	Horseweed/marestail ironweed lantana plantain
Vista XRT Remedy	9 fl oz/acre + 1 pt/acre	blackberry rose, multiflora	tropical soda apple wax myrtle persimmon

Spot Treatments

Spot treatments may be applied with a calibrated boom or with hand sprayers according to directions provided below.

Hand-Held Sprayers: Hand-held or backpack sprayers may be used for spot applications of Vista XRT if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on an area of 1,000 sq ft. The amount of Vista XRT (fl oz or ml) in the table should be mixed with 1 gallon or more of water and applied to an area of 1,000 sq ft. To calculate the amount of product required for larger areas, multiply the table value (fl oz or ml) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3,500 sq ft, multiply the table value by 3.5 (Calculation: $3,500 \div 1,000 = 3.5$). An area of 1000 sq ft is approximately 10.5 X 10.5 yards in size.

Amount of Vista XRT to Equal Specified Broadcast Rate (Mix with 1 Gallon or More of Water and Apply to 1,000 sq ft)				
6 fl oz/acre	9 fl oz/acre	12 fl oz/acre	17 fl oz/acre	22 fl oz/acre
0.14 fl oz (4.1 ml)	0.21 fl oz (6.2 ml)	0.28 fl oz (8.3 ml)	0.4 fl oz (11.7 ml)	0.56 fl oz (16.5 ml)

1 fl oz = 29.6 (30) ml

Restrictions:

- **Grazing and harvest restrictions:** There are no grazing restrictions for livestock, including lactating or non-lactating dairy animals. Meat animals must be withdrawn from treated forage at least 2 days before slaughter. Do not harvest grass for hay or silage from treated areas within 7 days of application.
- **Plant-back restriction:** Only forage grasses, wheat, barley, oats, field corn, sweet corn and grain sorghum may be planted in treated fields within 120 days following application of Vista XRT.
- **Vista XRT may injure or kill legumes.** Do not apply if the injury to legumes cannot be tolerated. Legumes may be less sensitive to herbicide injury after plant growth is mature and seed has set.
- **Maximum Application Rate:** Do not apply more than 22 fl oz of Vista XRT per acre per year.

®Trademark of Dow AgroSciences LLC

R362-004

EPA accepted: 12/10/07

Initial printing