

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

GoalTender[®] EPA Reg. No. 62719-447

Conifer Transplants and Container Stock (includes 2-0 Seedling) and Christmas Trees

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for GoalTender[®] herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of GoalTender according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for GoalTender.

Directions for Use

GoalTender[®] herbicide may be applied to conifer transplants and container stock (includes 2-0 seedling) and Christmas trees.

Agricultural Use Requirements: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 6 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Many container-grown, transplanted, and field grown conifers, including Christmas trees, are tolerant to preemergence and postemergence applications of GoalTender. Applied postemergence, GoalTender provides postemergence control of emerged weeds and preemergence residual control of many broadleaf weeds and grasses (see Key Weeds Controlled) at the beginning of this section.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	2 - 4	Transplanted and Container Grown Conifers and Christmas Trees: For best results, preemergence applications should be made immediately after transplanting seedlings or to weed-free container stock. Postemergence applications should be made to weeds less than 4 inches in height. Two applications may be necessary, in fall-transplanted conifer fields, for season-long weed control. The addition of a non-ionic surfactant (0.25% v/v) labeled for application to growing crops, enhances the activity of GoalTender on emerged weeds.

Precautions:

- Do not make over-the-top applications during periods of active conifer growth. Apply only before bud break or after new terminal growth has hardened off.

Crop-Specific Restrictions:

- Do not apply more than 4 pints of GoalTender per acre in a single application or more than 8 pints per acre per year.

In addition to those conifer species listed under the Conifer Seedbed section, the following conifer species have been shown to be tolerant to GoalTender:

Arborvitae	<i>Thuja occidentalis</i> <i>Thuja orientalis</i>
Juniper	<i>Juniperus chinensis</i> <i>Juniperus horizontalis</i> <i>Juniperus procumbens</i> <i>Juniperus sabina</i> <i>Juniperus scopulorum</i>
Red cedar	<i>Juniperus virginiana</i>
Western Hemlock	<i>Tsuga heterophylla</i>
Yew	<i>Taxus species</i>

®Trademark of Dow AgroSciences LLC
R204-064
EPA accepted: 05/28/09
Initial printing.