

Bayer Environmental Science

Bayer Environmental Science
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-800-331-2867

CoreTect™ Tree and Shrub Tablets Insecticide

For use on citrus and ornamentals to control
Asian citrus psyllid

EPA Reg. No. 432-1457

SLN No. CA-080016

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

It is a violation of federal law to use this product in a manner inconsistent with its label. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered Section 3 label.

THIS LABEL IS VALID UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

DIRECTIONS FOR USE

For use in and around residential areas, industrial and commercial buildings.

CITRUS: Crops of group 10 including: calamondin, *Citrus citron*, citrus hybrids (includes chironja, tangelo, and tango), grapefruit, kumquat, lemon, lime, mandarin (tangerine), pummelo, orange (sweet and sour), Satsuma mandarin, white sapote (*Casimiroa* spp.), and other cultivars and hybrids of these.

ORNAMENTALS: *Murraya* spp. (Curry Tree, Orange Jessamine)

IN-GROUND

Pest Controlled	CoreTect Tree and Shrub Tablets Insecticide
Asian citrus psyllid	2-3 tablets per inch of trunk diameter at breast height or per foot of shrub height

CONTAINERIZED

Pest Controlled	CoreTect Tree and Shrub Tablets Insecticide
Asian citrus psyllid	1 tablet/ft ³ of container media

Restrictions:

Do not apply more than 103 tablets per 10,000 ft² per year or 450 tablets per acre per year (.5 lb AI/Acre).

Preharvest interval (PHI): 0 days

Application Instructions:

CoreTect Tree and Shrub Tablets Insecticide is formulated to provide consistent delivery of active ingredient over time. Release of active ingredient is dependent on the presence of adequate soil moisture. For best results, irrigate the plant after treatment, and ensure that the plant receives regular irrigation for at least 2-3 months after treatment.

Typically, the fine absorptive roots of trees are located near the base of the tree, in the upper six inches of the soil. Apply CoreTect Tree and Shrub Insecticide Tablets near the base of the tree (i.e., within 6-24") to ensure rapid uptake by these absorptive roots. Space tablets evenly around the tree, 2-5" below the soil surface. Ensure that the tablets are completely covered with soil after the application.

For bushy trees and shrubs, numerous absorptive roots may be located near the drip-line. To apply CoreTect Tree and Shrub Tablets Insecticide to plants with a definite drip-line, place tablets 2-5" below the soil surface, spaced evenly around the tree and along the drip-line.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-800-331-2867

As with any plant-protection product, always read and follow label directions.