



SUPPLEMENTAL LABELING

AQUA STAR

42750-59

READ THE ENTIRE LABEL FOR AQUA STAR BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL. THIS SUPPLEMENTAL LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FORESTRY CONIFER AND HARDWOOD RELEASE

Directed Spray and Selective Equipment

This product may be applied as a directed spray or by using selective equipment in forestry conifer and hardwood sites, including Christmas tree plantations and silvicultural nurseries. Mix 2 to 6 quarts of a nonionic surfactant per 100 gallons of spray solution (0.5 to 1.5 percent by spray volume) for all spray applications. Use a surfactant with greater than 80 percent active ingredient.

In hardwood plantations, tank mixtures with Oust[®] may be used. In pine plantations, tank mixtures with Garlon[®] 4 or Arsenal[®] AC may be used. Comply with all site restrictions, forestry species limitations and precautions on the tank mix product label.

Avoid contact of spray, drift, mist or drips with foliage, green bark or non-woody surface roots of desirable species.

See all sections in the "APPLICATION EQUIPMENT AND TECHNIQUES" portion of this labeling for specific equipment recommendations and precautions.

For spray-to-wet applications, use a 1.5 percent spray solution for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 0.75 to 1.5 percent solution.

For low volume directed spray applications, use a 4 to 7.5 percent spray solution. Coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the unwanted vegetation is important.

For equipment calibrated for broadcast applications, use 1.5 to 7.5 quarts of this product per acre. Apply in 10 to 60 gallons of clean water per acre. Shielded application equipment may be used to avoid contact of the spray solution with desirable plants. Shields should be adjusted to prevent spray contact with the foliage or green bark of desirable vegetation.

Wiper application equipment may be used. See the "SELECTIVE EQUIPMENT" portion of this labeling for equipment and rate recommendations.

Broadcast Spray

Except where specifically recommended below, use only where conifers have been established for more than one year.

Application must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in the spring.

Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied. Damage can be accentuated if applications are made when conifers are actively growing, or are under stress from drought, flood water, improper planting, insects, animal damage or diseases.

This product may require use with a surfactant. Unless otherwise recommended in this section of this label, use Entry™ II surfactant at 10 to 30 fluid ounces per acre. Follow the instructions under the “MIXING” portion of this labeling.

For release of the following conifer species outside the Southeastern United States:

Douglas fir	Pines*
<i>Pseudotsuga menziesii</i>	<i>Pinus spp.</i>
Fir	Redwood, California**
<i>Abies spp.</i>	<i>Sequoia spp.</i>
Hemlock**	Spruce
<i>Tsuga spp.</i>	<i>Picea spp.</i>

*Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

**Use of a surfactant is not recommended for release of hemlock species or California redwood. In mixed conifer stands, injury to these species may result if a surfactant is used.

Apply 0.75 to 1.5 quarts of this product per acre as a broadcast spray.

Note: For release of Douglas fir with this product or recommended tank mixtures of this product, Entry™ II or a nonionic surfactant recommended for over-the-top foliar sprays may be used. To avoid possible conifer injury, Entry™ II rates should not exceed 20 fluid ounces per acre at elevations above 1500 feet, or 10 fluid ounces per acre in the coastal range or at elevations below 1500 feet in Washington and Oregon. Nonionic surfactants may be used at 2 fluid ounces per acre at elevations above 1500 feet, or 1 fluid ounce per acre in the coastal range or at elevations below 1500 feet. Use of surfactant rates exceeding those listed above may result in unacceptable conifer injury and are not recommended. Ensure that the nonionic surfactant has been adequately tested for Douglas fir safety before use. In Maine, up to 2.25 quarts per acre of this product or a tank mix with 1 oz/a of Arsenal® Applicators Concentrate may be used for the control of difficult species.

To release Douglas fir, pine and spruce species at the end of the first growing season (except in California), apply 0.75 to 1.125 quarts of this product per acre. Ensure that the conifers are well hardened off.

Oust® Tank Mixtures – To release jack pine, white pine and white spruce, apply 0.75 to 1.5 quarts of this product with 1 to 3 ounces (1 to 1.5 for white pine) of Oust® per acre. Make applications to actively growing weeds as a broadcast spray over the top of established conifers. Applications at these rates should be made after formation of conifer resting buds in the late summer or fall.

Arsenal® Applicators Concentrate Tank Mixtures – This product may be tank mixed with Arsenal® Applicators Concentrate for release of Douglas fir. Use 0.75 to 1.125 quarts of this product tank mixed with 2 to 6 fluid ounces of Arsenal® per acre. For release of balsam fir and red spruce, apply a mixture of 1.5 quarts of this product and 1 to 2.5 fluid ounces of Arsenal® Applicators Concentrate per acre. For release of the following conifer species in the Southeastern United States:

Eastern white pine	Shortleaf pine
<i>Pinus strobus</i>	<i>Pinus echinata</i>
Loblolly pine	Slash pine
<i>Pinus taeda</i>	<i>Pinus elliotii</i>
Longleaf pine	Virginia pine
<i>Pinus palustris</i>	<i>Pinus virginiana</i>

Apply 1.125 to 1.875 quarts of this product per acre as a broadcast spray during late summer or early fall after the conifers have hardened off. For applications at the end of the first growing season, use 0.75 quart per acre of this product alone or in a recommended tank mixture.

Arsenal[®] Applicators Concentrate Tank Mixtures – Apply 0.75 to 1.5 quarts of this product with 2 to 16 fluid ounces of Arsenal[®] Applicators Concentrate per acre as a broadcast spray for conifer release. Use only on conifer species that are labeled for over-the-top sprays for both products. Use the higher recommended rates for dense, tough-to-control woody brush and trees.

Read and carefully observe the label claims, cautionary statements and all information on the labels of each product used in these tank mixtures. Use according to the most restrictive precautionary statements for each product in the mixture.

Herbaceous Release

When applied as directed, this product plus listed residual herbicides provides postemergence control of the annual weeds and control or suppression of the perennial weeds listed in this label, and residual control of the weeds listed in the residual herbicide label. Make applications to actively growing weeds as a broadcast spray over the top of labeled conifers.

Oust[®] Tank Mixtures – To release loblolly pines, apply 12 to 18 fluid ounces of this product, plus 2 to 4 ounces of Oust[®] per acre. To release slash pines, apply 9 to 12 fluid ounces of this product, plus 2 to 4 ounces of Oust[®] per acre.

Mix up to 3.2 fluid ounces per acre of Entry[™] II with the recommended rate of this product plus Oust[®]. Applications can be made over newly planted pines after the emergence of herbaceous weeds in the spring or early summer. Best results are obtained from applications made in May and June.

Weed control may be reduced if water volumes exceed 25 gallons per acre for these treatments. Atrazine Tank Mixtures – To release Douglas fir, apply 0.75 quart of this product, plus 4 pounds active ingredient of atrazine per acre. Apply only over Douglas fir that has been established for at least one full growing season. Apply in the early Spring, usually mid-March through early April. Injury will occur if applications are made after bud swell in the Spring. Do not add surfactant to this mix for this use.

Always read and follow the manufacturer's label recommendations for all herbicides and surfactants used.

AD112907

Manufactured by:
Albaugh, Inc.
Ankeny, IA 50021