

SPECIAL LOCAL NEEDS REGISTRATION FOR NORTH DAKOTA ONLY

READ THE ENTIRE LABEL FOR GLY STAR™ PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. FOLLOW ALL DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

“Label” as used in this supplemental labeling refers to the label booklet for Gly Star™ Plus and this supplement.

Albaugh GLY STAR™ PLUS

EPA Reg. No. 42750-61
SLN No. ND-01-0006

FOR PREHARVEST APPLICATION IN FLAX IN THE STATE OF NORTH DAKOTA

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See “GENERAL INFORMATION”, “MIXING”, and “APPLICATION EQUIPMENT AND TECHNIQUES” sections of the label booklet for Gly Star™ Plus for essential product performance information.

RECOMMENDATIONS

This product may be applied as a spot treatment or a broadcast application to mature flax infested with Canada thistle or perennial sowthistle.

USE INSTRUCTIONS: This product may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of flax. Apply between 32 fluid ounces and 96 fluid ounces in 3 to 20 gallons of water per acre when the crop is physiologically mature and nearly ready to harvest. Apply when the crop is 30 percent or less grain moisture; approximately 80 percent of the seed should show a color change at this stage.

Either ground broadcast or aerial applications may be made. A 2.5 percent solution can be used when using hand-held or backpack sprayers.

PRECAUTIONS, RESTRICTIONS: Apply at least 7 days before harvest. Greater efficacy against perennial weeds may be achieved if a period of 10 to 14 days is allowed between treatment and crop harvest. Only one application per year may be made. Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Gly Star™ Plus herbicide label. Do not apply this product through any type of irrigation system. Do not add surfactants, additives containing surfactant, buffering agents or pH adjusting agents to this product. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

All other applicable instructions, restrictions, and precautions on the label booklet for this product must be followed, including the restricted reentry interval (REI) of 4 hours. Applicators and other handlers of this product for this use must wear the following personal protective equipment: long-sleeved shirt, long pants, shoes, socks.

Spray coverage should be uniform and complete, but do not spray weed foliage to the point of runoff. Do not apply when heavy rainfall is anticipated or when irrigation is scheduled soon after application. To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

Read the “CONDITIONS OF SALE AND WARRANTY” in the label booklet for Gly Star™ Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 500