



Section 2(ee) Recommendation

- Products: Bicep II MAGNUM[®] Herbicide, EPA Reg. No., 100-817
Bicep Lite II MAGNUM[®] Herbicide, EPA Reg. No., 100-827
Callisto[®] Herbicide, EPA Reg. No., 100-1131
- Use: Bicep II MAGNUM or Bicep Lite II MAGNUM in tank mixture with Callisto at 3 fl oz/Acre for improved postemergence control of selected annual broadleaf weeds in field corn grown, production seed corn and corn grown for silage
- States: Colorado, Iowa, Illinois, Indiana, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota, Maryland, Pennsylvania and Wisconsin

This recommendation is made under the authority of FIFRA Section 2(ee) and recommends Bicep II MAGNUM or Bicep Lite II MAGNUM in tank mixture with Callisto at 3 fl oz./Acre in field corn grown, production seed corn and corn grown for silage in the above noted states.

The user must refer to the federally approved labels for the above noted products and read and follow all directions for use, restrictions, and precautions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FOR POSTEMERGENCE AND RESIDUAL CONTROL BROADLEAF WEEDS AND CONTROL OR PARTIAL CONTROL OF ANNUAL GRASS WEED IN CORN:

Apply Callisto at 3.0 fl. oz. per acre in tank mixture with the recommended rate of Bicep II Magnum or Bicep Lite II Magnum. Apply in 10-30 gallons of water per acre. When weed foliage is dense, use a minimum of 20 gallons of water per acre. Flat fan nozzles are recommended for optimum postemergence coverage.

RESTRICTIONS

- 1) This tank mix will control most annual broadleaf weeds and provide control, or partial control, of some annual grass weeds as listed on the Callisto and Bicep II MAGNUM or Bicep Lite II MAGNUM labels (see individual product labels for a complete list of broadleaf weeds controlled and application timings).
- 2) This tank mix should not be applied to corn exceeding 12 inches in height.

- 3) Do not add UAN (28-0-0) or ammonium sulfate to the spray mixture or unacceptable crop injury may occur.
- 4) Temporary crop response may occur under certain weather conditions or if the crop is suffering from stress.
- 5) The user must have this recommendation in its possession at the time of use.

This recommendation is permitted under Section 2(ee) of FIFRA, as amended, and has not been submitted to or been approved by EPA.

All applicable directions, restrictions and precautions on the EPA-registered labels are to be followed.

This recommendation must be in the possession of the user at the time of pesticide application.

This FIFRA Section 2(ee) bulletin contains new or additional directions for use of this product which may not appear on the package label. Read and carefully observe precautionary statements plus all other information appearing on the product label.

Effective Date: February 25, 2002

Bicep II MAGNUM[®], Bicep Lite II MAGNUM[®], and Callisto[®] are registered trademarks of Syngenta.

Syngenta Crop Protection, Inc. (legacy company formed by the merger of Novartis Crop Protection, Inc. and Zeneca AG Products, Inc.)
Greensboro, NC 27409

CO1131016AC0202

COCallistoBicepIIMagSec2ee 2-25-02