

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF INDIANA.**



EPA Reg. No. 81880-6-10163

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

GLYPHOSATE-TOLERANT FIELD CORN ONLY

A tank mixture of YUKON Herbicide plus glyphosate may be used for Glyphosate Tolerant (GT) corn hybrids ONLY for enhanced knockdown and residual control of annual broadleaf weeds and nutsedge.

Yukon Use Rate Range: 2 - 8 ounces per acre.

Glyphosate Use Rate Range: 0.56 - 1.125 lbs. acid equivalence per acre.

If the glyphosate formulation selected does not contain a proprietary adjuvant system, then the addition of nonionic surfactant (NIS) is warranted at a rate of 0.25% v/v of spray mix. The addition of spray grade ammonium sulfate (AMS) at 8.5 - 17 lb/100 gal spray mix (or equivalency if a liquid AMS formulation is utilized) or 28% or 32% Urea Ammonium Nitrate (UAN) at 1-2% v/v is also required as an additive.

Corn Growth Stage: YUKON Herbicide plus glyphosate may be applied over-the-top or with drop nozzles to field corn from spike up to 30-36 inches tall dependent on the glyphosate formulation selected. Consult selected glyphosate product label for maximum crop stage for postemergence applications. The use of drop nozzles is recommended for applications made to GT corn over 20 inches in height to ensure proper coverage of weeds.

Following application to foliage, corn may be grazed or harvested for feed after the crop reaches the ensilage (milk) stage, at least 30 days after foliar application, unless further restricted by the glyphosate product label.