

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Lorsban[®] Advanced

EPA Reg. No. 62719-591

Special 2(ee) Recommendation[†]

(For Distribution and Use Only in the States of Oregon and Washington)

Control of Root Maggots, Cutworms, Symphylans, Billbugs, Wireworms and Grubs in Vegetables¹ Using Broadcast Application

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Lorsban Advanced according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

¹broccoli, Brussels sprout, cabbage, cauliflower, Chinese cabbage, collards, kale, kohlrabi, radish, rutabaga, turnip

Refer to product label for Lorsban Advanced for General Use Precautions, Mixing and Application instructions.

Preplant Incorporation Application for Direct Seeded or Transplanted Crops

Apply Lorsban Advanced as a broadcast spray to the soil surface using power-operated ground spray equipment. Use a total spray volume of 10 gallons per acre or more. On the day of treatment, incorporate Lorsban Advanced into the top 2 to 4 inches of soil using a disc, field cultivator, or equivalent equipment.

Crop	Target Pests	Lorsban Advanced (pints/acre)
cauliflower	billbugs cutworms grubs root maggot symphylans wireworms	4
broccoli		4.5
Brussels sprout		
cabbage		
Chinese cabbage		
collards		
kale		
kohlrabi		
turnip		
radish		5.5
rutabaga	4.5	

Specific Use Precautions:

Insecticides, including Lorsban Advanced, may contribute to the stress of plants under certain environmental conditions. This stress may reduce plant stand or interfere with normal plant development. Herbicides used preplant incorporated may interact with insecticides and enhance this stress.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 30 days of harvest;

- Do not foliarly apply any chlorpyrifos product labeled for foliar application within 10 days of a soil application of Lorsban Advanced.
- Do not make more than one soil application of any chlorpyrifos product per crop.
- Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours (3 days for cauliflower) unless PPE for early entry is worn.
- **Rutabaga:** Do not use rutabaga tops for food or feed purposes.

®Trademark of Dow AgroSciences LLC

R368-066

Issued: 03/30/10

Initial printing.