

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rodeo[®]

EPA Reg. No. 62719-324

2(ee) Recommendation[†]

For Distribution and Use Only in the State of South Carolina

Control or Suppression of Bamboo

ATTENTION

[†] This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to container for Rodeo[®] herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of Rodeo is subject to all use precautions and limitations imposed by the label affixed to the container for Rodeo.

Directions for Use

Rodeo may be used on roadside rights-of-way in South Carolina for control or suppression of bamboo. When applied as directed, Rodeo will control or partially control bamboo. Use a higher rate in the rate range for dense stands and larger plants.

Mow or cut and allow bamboo to resprout to have sufficient foliage for the spray solution to cover the foliage completely. Optimum control or suppression is achieved when applications are made between August and October (prior to frost). It is important to note that one application of Rodeo plus surfactant will not eradicate bamboo. Several mowing and spray applications are generally required for complete control of bamboo to be achieved.

Apply the specified rate (see below) plus a surfactant (typically 1/4 to 1/2% V/V), such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the Mixing and Application Instructions section of the product label and the surfactant manufacturer label for more information.

Where repeat applications are necessary, do not exceed 8 quarts per acre per year.

Method of Application	Application Rate	Spray Volume (gal/acre)
ground broadcast	1.5 - 7.5 qt/acre	10 to 60
handgun spray-to-wet	0.75 - 2%	spray-to-wet
handgun or backpack low volume directed spray	4% - 10%	spray to cover

[®]Trademark of Dow AgroSciences LLC

R148-005

Issued: 01/21/11

Initial printing.