

# Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## GoalTender<sup>®</sup>

EPA Reg. No. 62719-447

2(ee) Recommendation†

### For Use and Distribution in the State of Texas Only

#### Spray Volume for Use on Tree Fruit, Nut and Vine Crops (Dormant Application)

##### ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for GoalTender herbicide.

#### Directions for Use

##### Spray Volumes when Tank Mixing with Glyphosate and other contact herbicides

**Application Methods and Recommended Cultural Practices - Postemergence Weed Control:** Apply the specified rate in a broadcast spray volume of 20 or more gallons of water per acre (a minimum 10 gallons if applying GoalTender in tank mix with glyphosate or other contact herbicides which specify 10 gallons). Because GoalTender also has contact activity, complete and uniform coverage of weed foliage is essential for optimum postemergence control. Increase the spray volume to ensure complete and uniform coverage as weed height and density increases or in the presence of heavy weed or crop residue. Postemergence applications of GoalTender are most effective when made to weeds at the seedling stage. Applications made later than the 4-inch or 4 leaf stage may result in partial control or suppression. Postemergence applications should be made to seedling grasses not exceeding the 2-leaf stage. The addition of 0.25% v/v (2 pints per 100 gallons of spray) of a registered nonionic surfactant will enhance herbicidal effectiveness in controlling emerged weeds.

<sup>®</sup>Trademark of Dow AgroSciences LLC

R204-073  
Issued: 03/11/11  
Initial printing.