

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban[®] Advanced

EPA Reg. No. 62719-591

Special 2(ee) Recommendation[†]

(For Distribution and Use Only in the State of Florida)

Low Volume Application in Citrus Fruits¹

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Lorsban Advanced according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

¹Citrus fruits (crop group 10) including calamondin, chironja, citrus citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin (tangerine), pummelo, satsuma mandarin, sour orange, sweet orange, tangelo, tangor

Refer to product label for Lorsban Advanced for Use Precautions and Restrictions, Mixing and Application instructions.

Apply Lorsban Advanced at the rate of 3 to 5 pints per acre as a low volume application (e.g., 2 to 5 gallons of water) to control Asian citrus psyllid. Low volume applications should not be made when air speed is more than 5 mph. Regardless of the application method used (air, low volume, airblast), treat only a few acres when using the lower rate or a new application method to determine the effectiveness in the citrus grove.

When making low volume applications of Lorsban Advanced adjacent to permanent bodies of water such as rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds, the setback or buffer zone is 50 feet.

Do not make more than two applications of Lorsban Advanced or other product containing chlorpyrifos in a year. Read the label affixed to the container for Lorsban Advanced and follow all use directions, precautions, and restrictions for each crop.

[®]Trademark of Dow AgroSciences LLC

R368-072

Issued: 03/03/11

Initial printing.