
SUPPLEMENTAL LABEL



Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300

QUILT®

FUNGICIDE

Broad spectrum fungicide for control of plant diseases

Active Ingredient:	
Azoxystrobin.....	7.0%
Propiconazole	11.7%
Other Ingredients:	81.3%
Total:	100.0%

Contains 1.04 lbs. a.i. propiconazole and 0.62 lb. a.i. azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 100-1178

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Quilt as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cereals Wheat See next section for other cereals	Early season suppression of: Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	7 – 14	Apply Quilt in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control.
	Control of Leaf Diseases: Rust (<i>Puccinia</i> spp.) Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>) Helminthosporium Leaf Blight (<i>Drechslera tritici-repentis</i>) Spot Blotch (<i>Bipolaris sorokiniana</i>) Net Blotch (<i>Pyrenophora teres</i>) Kernel Blight (<i>Alternaria</i> spp.)	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Quilt is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl. oz./A may be applied. Applications may be made no closer than a 14-day interval. Quilt can be applied through full head emergence (Feekes growth stage 10.5). Do not apply after this stage to avoid possible illegal residues.
	Foot Rot/Eyespot (<i>Tapesia</i> spp.)	14	Apply full rate of Quilt plus half the rate recommended of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.

Application Instructions: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 2 applications/A/season.
- 2) Do not apply after Feekes 10.5.
- 3) Do not harvest within 30 days for forage or hay.
- 4) Do not apply more than 28.0 fl. oz./A/season of Quilt.
- 5) Do not apply more than 0.22 lb. a.i. propiconazole containing products/A/season.
- 6) Do not apply more than 0.40 lb. a.i. azoxystrobin containing products/A/season.
- 7) Under certain environmental conditions, tank mixes of Quilt plus herbicides and/or fertilizers may cause crop injury.

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cereals Barley Triticale	Early season suppression of: Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	7 – 14	Apply Quilt in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control.
	Control of Leaf Diseases: Rust (<i>Puccinia</i> spp.) Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>) Helminthosporium Leaf Blight (<i>Drechslera tritici-repentis</i>) Spot Blotch (<i>Bipolaris sorokiniana</i>) Barley Scald (<i>Rhynchosporium secalis</i>) Barley Stripe (<i>Pyrenophora graminea</i>) Net Blotch (<i>Pyrenophora teres</i>) Kernel Blight (<i>Alternaria</i> spp.)	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Quilt is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl. oz./A may be applied. Applications may be made no closer than a 14-day interval. Quilt can be applied up to 45 days prior to harvest.
	Foot Rot/Eyespot (<i>Tapesia</i> spp.)	14	Apply full rate of Quilt plus half the rate recommended of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.

Application Instructions: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 2 applications/A/season.
- 2) Do not apply within 45 days of harvest.
- 3) Do not harvest within 30 days for forage or hay.
- 4) Do not apply more than 28.0 fl. oz./A/season of Quilt.
- 5) Do not apply more than 0.22 lb. a.i. propiconazole containing products/A/season.
- 6) Do not apply more than 0.40 lb. a.i. azoxystrobin containing products/A/season.
- 7) Under certain environmental conditions, tank mixes of Quilt plus herbicides and/or fertilizers may cause crop injury.

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Corn Field Pop (Includes Seed Production)	Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium leaf blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>)	7 - 14	<p>Note: Applications prior to tasseling may impose stress on the plant that could inhibit proper kernel development, especially under environmentally stressful conditions.</p> <p>For leaf blights apply Quilt when disease first appears. Continue on a 7 to 14 day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.</p> <p>Apply no more than 2 applications of Quilt or any other Group 11 fungicide per year.</p>
	Anthracnose leaf blight (<i>Colletotrichum graminicola</i>) Rusts (<i>Puccinia</i> spp.) Gray Leafspot (<i>Cercospora zea-maydis</i>) Eye Spot (<i>Aureobasidium zeae</i>)	10.5 - 14	<p>Apply Quilt when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.</p> <p>Apply no more than 2 applications of Quilt or any other Group 11 fungicide per year.</p>

Application Instructions: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply within 30 days of harvest for forage, grain, or stover.
- 2) Do not apply more than 56 fl. oz./A/season of Quilt.
- 3) Do not apply more than 28 fl. oz. (0.224 lb. a.i. propiconazole) for field corn harvested for forage.
- 4) Do not apply more than 0.45 lb. a.i. propiconazole containing products/A/season.
- 5) Do not apply more than 2.0 lbs. a.i. azoxystrobin containing products/A/season.

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Corn, Sweet Sweet corn (Includes Seed Production)	Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium leaf blights (<i>H. maydis, H. turcicum, H. carbonum</i>)	7 - 14	For leaf blights apply Quilt when disease first appears. Continue on a 7 to 14 day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate. Alternate applications of Quilt with Tilt or another product with a different mode of action than Group 11 fungicides.
	Anthracnose leaf blight (<i>Colletotrichum graminicola</i>) Rusts (<i>Puccinia</i> spp.) Gray Leafspot (<i>Cercospora zea-maydis</i>) Eye Spot (<i>Aureobasidium zeae</i>)	10.5 - 14	Apply Quilt when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Alternate applications of Quilt with Tilt or another product with a different mode of action than Group 11 fungicides.

Application Instructions: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt may be applied by ground, air, or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl. oz./A/season of Quilt.
- 2) Do not apply to sweet corn within 14 days of harvest for ears or forage.
- 3) Do not apply more than 0.45 lb. a.i. propiconazole containing products/A/season.
- 4) Do not apply more than 2.0 lbs. a.i. azoxystrobin containing products/A/season.

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Sorghum	Anthracnose <i>(Colletotrichum graminicola)</i> Gray leaf spot <i>(Cercospora sorghi)</i> Ergot <i>(Claviceps sorghi)</i>	14	<p>For foliar diseases, Quilt applications should begin prior to disease development. Applications may be made by ground, air or chemigation.</p> <p>For ergot: Make the first application at or just prior to flowering. Repeat on a 5-7 day interval. Apply up to 4 times.</p> <p>Resistance Management: Do not apply more than two sequential applications of Quilt or other Group 11 containing fungicides before alternation with a fungicide that is not in Group 11.</p>

Application Instructions: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. Quilt may be applied by ground, air or chemigation.

Specific Use Restrictions:

- 1) Do not apply more than 55 oz of Quilt/A per season
- 2) Do not apply more than 0.22 lb propiconazole a.i./A (27.5 oz Quilt) on sorghum harvested for forage.
- 3) Do not apply more than 0.45 lb. a.i. propiconazole containing product/A/season.
- 4) For grain and stover, do not apply more than 0.75 lbs. a.i./A per season of azoxystrobin-containing products.
- 5) For forage, do not apply more than 0.5 lbs. a.i./A per season of azoxystrobin-containing products.
- 6) Do not graze livestock or cut for green chop or silage within 30 days of application.
- 7) Do not apply within 30 days of harvest for forage
- 8) Do not apply within 21 days of harvest for grain and stover.

Manufactured for:
 Syngenta Crop Protection, Inc.
 P.O. Box 18300
 Greensboro, North Carolina 27419-8300

SCP 1178A-S10 0209

QUILT 1178A-S10 0209 - bb - 3-5-09