

---

**PRODUCT INFORMATION**

---



**Syngenta Crop Protection, Inc.**  
Greensboro, NC 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

**SUPPLEMENTAL LABEL FOR QUILT™**

**FUNGICIDE**

**EPA Reg. No. 100-1178**

Grasses grown for seed removal of feeding restriction

Active Ingredient:	
Azoxystrobin .....	7.0%
Propiconazole .....	11.7%
Other Ingredients:	81.3%
Total:	100.0%

Contains 1.04 lbs. a.i. propiconazole and 0.62 lb. a.i. azoxystrobin per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

**Before using Quilt as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**DIRECTIONS FOR APPLICATIONS**

Crop	Target Diseases	Use Rate fl oz./product/A	Remarks
<b>Grasses</b> (grown for seed) (Nebraska, Oregon, Washington, Idaho, and Minnesota only)	Rusts ( <i>Puccinia</i> spp.) Powdery mildew ( <i>Erysiphe graminis</i> ) Selenophoma stem eyespot ( <i>Selenophoma</i> spp.) Ergot Stem Diseases	14-27.5	Apply Quilt when powdery mildew infections, <i>Selenophoma</i> infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. To maximize control of severe rust pressure, apply 27.5 fl. oz./A (except bluegrass apply 14 fl. oz./A) and make applications at 14-day intervals until the seed is mature. Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin application early in the growing season.  Make no more than 2 sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.  Apply Quilt in a minimum of 20 gals. of water per acre for ground application, or in a minimum of 10 gals. of water per acre for aerial application. Quilt may also be applied by chemigation.
<p><b>Specific Use Restrictions:</b> Do not feed hay cut within 20 days of the last application.                      Do not graze treated areas within 140 days of the last application.                      Do not apply more than 110 fl. oz./A/season of Quilt.                      Do not apply more than 0.90 lb. a.i. propiconazole containing products/A/year.                      Do not apply more than 0.75 lb. a.i. azoxystrobin containing products/A/year.</p>			

Trademark of Syngenta

©2004 Syngenta

Syngenta Crop Protection, Inc.  
 Greensboro, North Carolina 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

SCP 1178A-S1 0304

QUI 1178A-S1 0304 - nsp - 3-29-04