
PRODUCT INFORMATION



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com

SUPPLEMENTAL LABEL FOR SEQUENCE™ HERBICIDE FOR AERIAL APPLICATION IN CALIFORNIA ONLY

Active Ingredient:

*Glyphosate: N-(phosphonomethyl) glycine	21.8%
**S-metolachlor (CAS No. 87392-12-9)	29.0%
<i>Other Ingredients:</i>	49.2%
<i>Total:</i>	100.0%

*Contains 2.25 pounds of glyphosate acid per U.S. gallon.

**Contains 3 pounds of S-metolachlor per U.S. gallon.

EPA Reg. No. 100-1185

KEEP OUT OF REACH OF CHILDREN.

CAUTION

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Sequence Herbicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling supersedes the Directions for Use on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SCP 1185-S1 0804

DIRECTIONS FOR USE

It is a violation of federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

See "GENERAL INFORMATION" and "DIRECTIONS FOR USE" sections of the label booklet for essential product information.

See the "CROPS" section of the label booklet for specific recommendations on the use of this product.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THE SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL GLYPHOSATE TOLERANT CROPS), PLANTS, TREES OR OTHER DESIRABLE VEGETATION, BECAUSE SEVERE DAMAGE OR DEATH OF PLANTS MAY RESULT.

Aerial applications of this product are allowed in the following situations:

1. In fallow and reduced tillage systems prior to the emergence or of labeled crops.
2. Over-the-top applications to glyphosate tolerant corn and to glyphosate tolerant cotton (including Roundup Ready® corn and cotton). Refer to the Sequence Herbicide label for directions for postemergence application to glyphosate tolerant corn and cotton.

Do not replant treated areas to crops other than those listed in the label booklet for 30 days following application. Do not exceed dosage limitations for the replant crop.

When applied as recommended under the conditions described, this product controls annual and perennial weeds listed in the label booklet.

For aerial application of Sequence in fallow and reduced tillage culture of crops, in tank mixture with 2,4-D; only use the amine salt of 2,4-D.

DO NOT EXCEED A MAXIMUM RATE OF 64 FLUID OUNCES PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN FALLOW AND REDUCED TILLAGE CULTURE.

DO NOT EXCEED A MAXIMUM RATE OF 34 FLUID OUNCES PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN CORN. THIS RESTRICTION ALSO APPLIES TO OVER-THE-TOP APPLICATIONS IN GLYPHOSATE TOLERANT CORN AND GLYPHOSATE TOLERANT COTTON.

Aerial Equipment

Use the recommended rates of this product in 3 to 15 gallons of water per acre. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

AVOID DRIFT – DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the air-stream and do not increase spray volume by increasing nozzle pressure. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application – To avoid streaking, uneven, or overlapped application, use appropriate marking devices. Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

Read the “Conditions of Sale and Limitation of Warranty and Liability” in the label for Sequence Herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Sequence™ and the Syngenta logo trademarks of a Syngenta Group Company

Roundup Ready® trademark of Monsanto Company

©2005 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com