
SUPPLEMENTAL LABELING

Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1212A-S1 0809

GROUP	9	HERBICIDE
-------	----------	-----------

Touchdown CT
Herbicide

Nonselective Foliar Systemic Herbicide for Weed Control

Active Ingredient:

*Glyphosate, N-(phosphonomethyl) glycine 36.5%

Other Ingredients: 63.5%

Total: 100.0%

*Contains 4.17 pounds of glyphosate acid in each gallon, in the potassium salt form.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1212

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Touchdown CT as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

VEGETABLE CROPS

Touchdown CT may be used on the vegetable crops listed below:

Amaranth	Ground Cherry
Arrugula	Guar
Artichoke (Jerusalem)	Horseradish
Asparagus	Kale
Beans (all)	Kohlrabi
Beet, garden	Leeks
Beet, sugar	Lentils
Broccoli (all)	Lettuce
Brussels Sprouts	Melons (all including citron, crenshaw, honey balls, honeydew, mango, musk, Persian)
Cabbage (all)	Mizuna
Cabbage (Chinese)	Mustard greens
Cantaloupe	Okra
Cardoon	Onions (green, spring, Japanese bunching)
Carrot	Parsley, turnip-rooted
Casaba	Parsnip
Cavalo Broccolo	Peas (all)
Cauliflower	Pepinos
Celeriac	Pepper (all)
Celery	Potato (Irish)
Celery (Chinese)	Pumpkin
Celtuce	Purslane
Chard (Swiss)	Radish
Chayote	Radish, oriental (daikon)
Chervil	Rape greens
Chick peas	Rhubarb
Chicory	Rutabaga
Chrysanthemum	Salsify, black
Collards	Salsify (oyster plant)
Corn salad	Salsify (Spanish)
Cress	Shallots
Cucumber	Spinach (all)
Dandelion	Spinach, mustard
Dock (sorrel)	Squash (summer, winter)
Eggplant	Sweet potato
Endive	Tomatillo
Fennel (Florence)	Tomato
Garlic	Turnip
Gherkin	Watercress
Ginseng	Watermelon
Gourd, edible	Yams

Method of Application: Broadcast application before transplanting or before, during, or after planting but prior to crop emergence if direct seeded; spot spray; wiper/wick (rutabaga only); postharvest. In addition, preharvest or spot application in dry beans, peas, lentils and chickpeas.

Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES,** and **APPLICATION EQUIPMENT AND TECHNIQUES** sections of this label. Refer to **WEEDS CONTROLLED** for application rates and timing.

Use Precautions and Restrictions for Vegetable Crops

- For **Asparagus**, do not apply broadcast within one week of emergence of first spears. Spot applications can be made immediately after cutting, but before emergence. Spears can be harvested 5 days after spot application. Postharvest applications can be made as a directed or shielded spray avoiding contact of the spray with the ferns, stems, or spears.
- Wiper/wick applications to rutabagas must be made at least 14 days before harvest.
- Make one preharvest or one spot application in **dry beans** with no more than 24 fl oz/A or in **peas, lentils, and chickpeas** with no more than 69 fl oz/A at least 7 days before harvest.
- Employ at least a 30 day plant back interval between treatment and replanting for any crop not listed on this label.
- Do not feed treated vines and hay to livestock.
- Do not combine a preharvest with a spot application on the same crop area.
- Do not treat cowpeas or field (feed) peas, these crops are considered as livestock feed.
- Wait 3 days after application before planting cantaloupe, casaba, chayote, Chinese okra, Chinese waxgourd, cucumber, cucuzza, edible gourd, eggplant, gherkin, gourds, groundcherry, melons (all), pepper (all), pumpkin, squash, tomatillo, watercress, and watermelon.
- If transplanting into plastic mulch, ensure residues of this product are removed from the plastic prior to transplanting. Residues can be removed by a minimum of ½ inch of sprinkler irrigation or rainfall.
- Preharvest application is not recommended for legumes grown for seed as reduction in germination or vigor may occur.

Tank Mixtures With Residual Herbicides for Preplant/Preemergence Use in Vegetables

Touchdown CT can be tank mixed with the following herbicides for control or suppression of annual and perennial weeds provided that the tank mix product label allows use of the product. Refer to the **WEEDS CONTROLLED** section for application rates and timing. Apply Touchdown CT at 0.7 to 5.8 pt/A in these tank mixes for control or suppression of annual and perennial weeds. For control or suppression of dense populations of weeds greater than 12 inches in height or weeds under stress, consider use rates at the higher end of the rate range.

Devrinol®
Command®
Dual MAGNUM®
Dual II MAGNUM®
Fusilade® DX
Goal®
Kerb®
Lorox®

Matrix®
Prefar®
Prowl®
Sencor®
Trifluralin
Treflan®
Turbo

Refer to individual product labels for precautionary statements, restrictions, rates, approved uses, and a list of weeds controlled.

Preharvest

Touchdown CT may be applied over-the-top of dry beans, peas, lentils and chickpeas to control weeds before harvest. Apply no more than 24 fl oz/A in dry beans or no more than 69 fl oz/A in peas, lentils, and chickpeas in 3-20 gallons of water per acre at the hard dough stage of the legume seed (30% or less grain moisture). Only one application per year may be made.

Spot Treatment

Touchdown CT may be applied as a spot treatment to control troublesome weeds in dry beans at no more than 24 fl oz/A, and in peas, lentils, and chickpeas at no more than 69 fl oz/A. Apply in 10-20 gallons of water using a ground sprayer or use a 2% solution in a hand-held sprayer. For best results, make applications at or beyond the bud stage of growth. Only one application per year may be made.

Dual Magnum®, Dual II Magnum®, Fusilade® DX, and the Syngenta logo trademarks of a Syngenta Group Company

Command® trademark of FMC Corporation

Devrinol® trademark of United Phosphorous, Inc.

Goal®, Kerb®, Treflan® trademarks of Dow AgroSciences

Lorox®, Matrix® trademarks of E. I. DuPont de Nemours & Co., Inc.

Prefar® trademark of Gowan LLC

Prowl® trademark of BASF Ag Products

Sencor® trademark of Bayer CropScience

©2010 Syngenta