

---

**SUPPLEMENTAL LABELING**

---

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

**Syngenta Crop Protection, LLC**

P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1431A-S4 1211**

**Gramoxone® SL 2.0**

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide  
Potato – (late blight disease control) For Use Only in the State of Maine

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride) ..... 30.1%

Other Ingredients: ..... 69.9%

Total: ..... 100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon.  
Contains alerting agent (odor), emetic, dye.

**EPA Reg. No. 100-1431**

**KEEP OUT OF REACH OF CHILDREN.**



**DANGER / POISON  
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- **NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.**
- **IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.**
- **DO NOT USE OR STORE IN OR AROUND THE HOME.**
- **DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.**
- **THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.**

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL 2.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**syngenta**

## DIRECTIONS FOR USE

Crop	Maximum Number of Applications	Use Pattern	Gramoxone SL 2.0 Rate/Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval	Precautions, Restrictions and Comments
<b>POTATO</b>  (late blight disease control)  For Use Only in the State of Maine	2	Broadcast  for control of late blight disease	2.5 pts.	Ground: 20 gals.	-	<ul style="list-style-type: none"> <li>• For Desiccation of Potato Plants to Control Late Blight Disease</li> <li>• <b>This is a CROP DESTROY use. Crop must be destroyed and tubers not allowed to enter channels of trade. All tubers must be disked-in after desiccation.</b></li> <li>• Apply broadcast over-the-top of infected plants.</li> <li>• Add a nonionic surfactant at 1.0 quart per 100 gallons of spray mix.</li> <li>• Make a second application if necessary to obtain additional desiccation where vine growth is dense.</li> <li>• Allow a minimum of 5 days between applications.</li> <li>• Do not graze livestock in treated area.</li> <li>• Do not apply this product through any type of irrigation system.</li> </ul>

Gramoxone® SL 2.0 and the Syngenta logo trademark of a Syngenta Group Company

©2012 Syngenta

GRAMOXONE SL 2.0 1431A-S4 1211 – tb – 02/08/12