

---

**RESTRICTED USE PESTICIDE  
Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision,  
and only for those uses covered by the Certified Applicator's certification.

---

**INTRA**

**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ALABAMA**

**GRAMOXONE® SL 2.0**

**EPA REG. NO. 100-1431  
EPA SLN NO. AL-120003**

**KEEP OUT OF REACH OF CHILDREN**

**Danger/Peligro  
POISON**



**For Suppression and/or Control of Palmer Amaranth in Peanut**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL  
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

---

**DIRECTIONS FOR USE**

---

It is a violation of federal law to use this product in a manner inconsistent with all applicable directions, restrictions, and precautions found in the labeling of the federally registered product and accompanying supplemental labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Gramoxone SL 2.0 label.

This label must be in the possession of the user at the time of application.

## Application Method

Gramoxone SL 2.0 can be applied through recirculating rope wick or carpet roller equipment for suppression or control of glyphosate-resistant Palmer amaranth or to prevent seed production. Best results have been achieved using carpet roller equipment.

Make application at least 6 inches above the peanut canopy, and set application equipment to avoid dripping. Low ground speed enhances coverage and hence weed control. Speeds less than 5 mph are recommended.

Suppression or control of large weeds is enhanced if application is made in late afternoon or early evening.

Follow application equipment manufacturer's instructions.

## Mixing Instructions and Application Rate

Mix 1 part of Gramoxone SL 2.0 with 1 part of water to prepare a 50% solution. Add nonionic surfactant containing 75% or more surface-active agent at a minimum of 0.25% v/v (2 pt/100 gal or 0.32 fl oz/gal) of the finished volume.

Adjust equipment to apply up to 2 pt/A of the herbicide-water mixture.

Follow all safety precautions on label when mixing and applying Gramoxone SL 2.0 and when cleaning application equipment.

## Restrictions

1. Do not apply more than 1 pt/A of Gramoxone SL 2.0 using the recirculating rope wick or carpet roller application method. If Gramoxone SL 2.0 was used as a burndown in no-till at a rate greater than 1 pt/A, then do not make this application.
2. Do not exceed 0.5 lb ai/A (1 qt/A) total for the entire growing season from all applications on this or any other supplemental or Section 3 Federal label for peanut
3. Make applications of Gramoxone SL 2.0 as early as possible for best result, but not less than 30 days before harvest to suppress weeds that may interfere with harvesting operations.
4. Do not allow livestock to graze in treated areas.
5. Do not feed hay or threshings from treated fields to livestock.

Section 24(c) registrant:  
Syngenta Crop Protection, LLC  
Greensboro, NC 27419-8300

Gramoxone® trademark of a Syngenta Group Company

AL1431098BA0612