



# INTIRA

## Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TENNESSEE

**Dual Magnum Herbicide  
For Weed Control in Beans  
(Soybean, Immature Seed; Snap Bean; Lima Bean; Southern Pea)**

EPA Reg. No. 100-816  
EPA SLN No. TN-160001

**This label expires and must not be distributed or used in accordance with this SLN registration after February 22, 2021**

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

---

### **BEANS (Soybean, Immature Seed; Snap Bean; Lima Bean; Southern Pea)**

---

#### **Postemergence Broadcast or Postemergence Directed Application**

Apply Dual Magnum at 0.5 to 0.75 pt/A after the first trifoliate stage of plant growth.

Application to plants with less than one trifoliate can result in unacceptable crop injury. When applied broadcast over the top, crop injury in the form of leaf spotting and speckling may be observed.

If a preemergence application of Dual Magnum was made to the crop, a postemergence application may be made provided the total amount of Dual Magnum applied does not exceed 2.0 pt/A. Refer to the Dual Magnum label for preplant and preemergence application directions.

#### ***Precautions:***

- 1) Application to wet plants or when conditions are extremely hot or humid will result in increased risk of crop injury.
- 2) The risk of crop injury is greater on lighter textured soils and with higher use rates, especially when coupled with heavy rains or when excessive multiple irrigations occur within 5 days of application.

- 3) Dual Magnum will not control emerged weeds; thus, other control tactics that provide weed control prior to Dual Magnum being activated are required.
- 4) All bean and pea cultivars have not been tested for tolerance. Experiment on a limited basis until on-farm confidence in these use patterns and rates are obtained.
- 5) Dual Magnum should be applied only with water as the carrier.

**Restrictions:**

- 1) Do not apply more than 0.75 pt/A of Dual Magnum postemergence.
  - 2) Do not apply more than 2.0 pt/A of Dual Magnum total per season.
  - 3) Do not apply to beans from the time of soil cracking until first trifoliolate.
  - 4) Do not harvest crops within 50 days of the Dual Magnum application.
  - 5) Do not feed animals with plant material treated with a post-emergence application of Dual Magnum.
- 

24(c) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, NC 27419-8300

Label Code: TN0816028AA0216