

---

**RESTRICTED USE PESTICIDE  
Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision,  
and only for those uses covered by the Certified Applicator's certification.

---

**syngenta**

**FIIRA**

**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS**

**GRAMOXONE® SL 2.0**

**Harvest Aid Applications on Cotton in Texas  
(except in the counties of Starr, Hidalgo, Willacy, and Cameron).**

**EPA REG. NO. 100-1431**

**EPA SLN NO. TX-120010**

This label expires and must not be distributed or used in accordance with this SLN registration  
after December 31, 2020.

**KEEP OUT OF REACH OF CHILDREN**

**DANGER/PELIGRO  
POISON**



**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL  
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

## DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

### Specific Crop Use Directions

Three multiple broadcast harvest aid applications of Gramoxone SL 2.0 may be used. **Do not exceed a total of 3 pints per acre per crop season.** Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.

Crop	Rate Per Acre	Precautions, Restrictions and Comments
Tankmixes with phosphate or chlorate defoliant	8.1 – 16.1 fl oz. alone <u>or</u> tank mixed with 1 1/2 - 2 gal. chlorate, <u>or</u> 1 1/2 - 2 pts. phosphate	<ul style="list-style-type: none"> <li>• Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.</li> <li>• In desert cotton areas or on rank vigorous cotton, use the 16.1 fl oz rate of Gramoxone SL 2.0</li> <li>• Apply when 75% or more of the bolls are open and remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited.</li> <li>• Follow Harvest Aid Use Precautions listed on the Federal label.</li> </ul>
Tankmixes for boll opening and defoliation of southern cotton	4.1 to 5.0 fl. oz.	<ul style="list-style-type: none"> <li>• To aid in defoliation and opening of mature bolls, Gramoxone SL 2.0 may be tank mixed with the following products: DEF<sup>®</sup> defoliant, DROPP<sup>®</sup> defoliant, PREP<sup>™</sup> PGR, FOLEX<sup>®</sup> defoliant</li> <li>• Apply when 60% or more of the bolls are open and the remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited.</li> <li>• Refer to tank mix product label for rate, directions, limitations and caution.</li> </ul>
Harvest aid for defoliation and boll opening of stripper or spindle harvested cotton	3.1 to 11.2 fl. oz.	<ul style="list-style-type: none"> <li>• Apply when 75% of the bolls are open and the remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity.</li> <li>• Gramoxone SL 2.0 may be applied alone or tank mixed with the cotton harvest aid products listed above.</li> <li>• To avoid leaf sticking, apply Gramoxone SL 2.0 as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14 days before harvest.</li> <li>• Cooler temperatures may cause a longer waiting period between application of Gramoxone SL 2.0 as a desiccant and defoliation/conditioning.</li> <li>• Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation.</li> </ul>
Harvest aid desiccation of stripper cotton or spindle harvested cotton late season	16.8 to 31.5 fl. oz.	<ul style="list-style-type: none"> <li>• Use to desiccate regrowth occurring after defoliation or desiccation.</li> <li>• Apply when 85% of the bolls are open and the remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity.</li> <li>• To avoid leaf sticking, apply Gramoxone SL 2.0 as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14 days before harvest.</li> </ul>

Crop	Rate Per Acre	Precautions, Restrictions and Comments
		<ul style="list-style-type: none"> <li>Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation.</li> </ul>
Desiccation of Regrowth	16.8 to 31.5 fl. oz.	<ul style="list-style-type: none"> <li>Use to desiccate regrowth after defoliation or desiccation.</li> <li>Regrowth is difficult to control, therefore, thorough coverage with the full recommended rate is necessary.</li> <li>Control is dependent upon growing condition and desiccation of small new growth may not always be complete.</li> <li>Use higher rate if regrowth is excessive.</li> </ul>

Refer to the Gramoxone SL 2.0 federal label for additional cotton recommended uses and nonionic surfactant recommendations.

**Additional Precautions, Restrictions and Comments:**

- It is advisable, because of extremes in environmental and plant conditions, to apply the range of rates on a small block of cotton to determine the rate that best fits your needs.
- Do not pasture lactating dairy animals.
- Do not pasture livestock in treated fields or feed treated foliage.
- When combined with phosphate defoliant, observe livestock cautions listed on the phosphate label.
- Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.
- Do not apply this product through any type of irrigation system.

Section 24(c) registrant:  
 Syngenta Crop Protection, LLC  
 P.O. Box 18300  
 Greensboro, NC 27419-8300

© 2015 Syngenta

Gramoxone® trademark of a Syngenta Group Company

Def®, Dropp®, Prep™ PGR, trademarks of Bayer CropScience  
 Folex® trademark of Amvac Chemical Corporation

Label Code: TX1431107BA0915