

# **TURF - SPRING, SUMMER AND FALL DISEASES - LARGE PATCH**

## General Information

### GENERAL USE DIRECTIONS

Quali-Pro Ipro 2SE is a fungicide recommended for control of spring, summer, fall and winter diseases on non-residential turf located on golf courses, sod farms and institutional areas where fine turf is grown, and on nonresidential ornamental flowers and foliage plants grown in field and landscape nurseries or greenhouses, and in conifer nurseries.

Follow all directions, recommendations and precautions and limitations on the label to obtain optimum disease control and to prevent crop injury.

DO NOT apply this product to residential turf and/or ornamentals.

DO NOT apply this product when the wind direction is toward aquatic areas. Refer to the Spray Drift management section and tank mix information in the label.

### Limitations, Restrictions, and Exceptions

#### TURF - SPRING, SUMMER AND FALL DISEASES - LARGE PATCH

Make first application in fall when conditions are favorable for disease development but no symptoms are visible. Apply in the spring as needed on 14-21 day spray intervals.

Not registered for use in California.

Applied according to good turf management practices, Quali-Pro Ipro 2SE is an effective tool in controlling spring, summer and fall diseases and in controlling certain winter diseases.

Mixing and Application Instructions: Use the tables in the label to determine the recommended rates depending on the disease. Prepare only the amount of spray mixture required for a single day's (12 hours) application to avoid decomposition of the product. Add the required amount of Quali- Pro Ipro 2SE to 0.5 to 10 gal water per 1,000 sq. ft. Agitate the spray mixture during mixing and application in a

properly calibrated sprayer. Apply when disease first appears or when disease development can be expected. Apply to ensure uniform coverage of the turf but do not drench.

Use the higher rate and/or shorter spray interval between applications for severe disease conditions. The lower rates and longer intervals between applications are recommended for light to moderate disease pressure.

Restrictions:

- DO NOT apply to turf cut higher than 1" on golf holes where water bodies are present.
- Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.
- DO NOT mow or irrigate treated areas until the grass is completely dry, which typically takes 24 hours post-treatment.
- DO NOT mix with any sticker, extender, or wetting agents.
- DO NOT allow animals to graze on treated turf and DO NOT feed clippings from treated turf to livestock or poultry.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[When conditions are favorable for disease development but no symptoms are visible.](#)

[In the spring as needed.](#)