

# **TURF (ESTABLISHED) - MODERATELY TOLERANT - BROADCAST APPLICATION**

## General Information

### GENERAL INFORMATION

EJECT 75DF herbicide may be applied postemergence to residential and nonresidential turfgrasses (Refer to Table 1 in the label) for the control of many broadleaf and grass weeds. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms.

EJECT 75DF herbicide is absorbed by foliage and roots and translocated throughout the plant. The control symptoms exhibited by broadleaf weeds include leaf and stem curl or twisting, and chlorosis. Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death. Refer to Tables 1, 2 and 3 in the label for turfgrass tolerance and susceptible weed species.

### APPLICATION INFORMATION

Apply EJECT 75DF herbicide to actively growing weeds as postemergence broadcast or spot sprays using the rate, turf species and growth stage indicated in Tables 1, 2 and 3 of the label. Do not exceed the recommended application rate or fail to comply with use restrictions listed in General Restrictions and Limitations.

To achieve consistent weed control, methylated seed oil is recommended. Refer to Tables 2 and 3 in the label for rates.

Adding adjuvants may cause slight leaf burn, but new growth is normal and turf vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. Additional stress from low mowing heights may also increase the possibility of turf injury. Chelated iron or sprayable soluble nitrogen fertilizer will reduce a slight yellowing that may occur on some turfgrass species.

Spot Applications:

Postemergence spot applications may be made to susceptible weeds in turfgrass that is tolerant to EJECT 75DF herbicide (see Tables 1 and 2 in the label). Apply 0.367 ounce of EJECT 75DF herbicide per 1,000 square feet of treated area. Spray coverage should be uniform and complete.

#### Mowing Information:

It is recommended to not mow 2 days before or after applying EJECT 75DF herbicide to maximize weed control and minimize potential turf injury. Clippings from the first three mowings should be left on the treated area.

#### Irrigation and Rainfall:

If soil moisture is not sufficient prior to EJECT 75DF herbicide application, irrigation may improve weed control. For best results, do not water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is recommended.

#### Extended Grass Control:

To extend grass control EJECT 75DF herbicide can be tank-mixed with Pendulum 3.3 EC, PRE-M 3.3 turf herbicide or Pendulum 60 DG to provide residual control of annual grasses. Consult the respective tank-mix labels for additional weeds controlled.

#### Seeding/Overseeding/Sprigging:

The use of EJECT 75DF herbicide before or after seeding or overseeding a turf area will not significantly interfere with the turfgrass seed germination and growth of those grass types identified as tolerant in Table 1. Consult Table 4 in the label for timing of applications concerning any seeding, overseeding or sprigging situation.

#### GENERAL RESTRICTIONS AND LIMITATIONS

- DO NOT apply more than 2 pounds of EJECT 75DF herbicide per acre (or 0.73 ounces per 1,000 square feet) in one year.
- DO NOT apply to golf course collars or greens.

- DO NOT apply to fine fescue unless it is part of a seed blend.
- DO NOT apply to bahiagrass, carpetgrass, St. Augustinegrass, centipedegrass, dichondra, or lawns or turf where desirable clovers are present.
- DO NOT apply within 4 weeks after seedling emergence of Kentucky bluegrass, creeping bentgrass, fine fescue blends and perennial ryegrass.
- DO NOT apply to exposed feeder roots of trees or ornamentals. Be particularly careful within the drip line of trees and other ornamental species.
- DO NOT apply into any ornamental bed.
- DO NOT use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Apply when wind speed is less than 10 mph as drift may cause damage or death of non-target area vegetation. DO NOT apply when conditions favor drift from target area.
- Use a lawn type sprayer with coarse spray as wind drift is less likely.
- Avoid mist and spray onto vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants, especially plants belonging to the Solanaceae family such as tomatoes, eggplants, and bell peppers.
- DO NOT pour spray solutions near these plants.
- DO NOT apply through any type of irrigation system.
- DO NOT use to formulate or reformulate any other pesticide product which is not registered by the EPA.
- For best results, weeds should not be under stress from lack of water, excessive water, low fertility, mowing shock, excessive hot or cold temperatures, or injury from other herbicide applications.

Limitations, Restrictions, and Exceptions

TURF (ESTABLISHED) - MODERATELY TOLERANT - BROADCAST APPLICATION

Additive Rate: Apply: 1.5 pints per acre (0.55 oz. per 1,000 sq. ft.) methylated seed oil.

For Creeping Bentgrass and Hybrid Bermudagrass: Yellowing that occurs on these species can be reduced by the addition of chelated iron or sprayable soluble nitrogen fertilizer (See Application Information, Additives in the label).

For Fescue, Fine: Apply EJECT 75DF herbicide to fine fescue only when it is part of a blend.

- Do not use on golf course greens and collars.
- Consult Table 4, Seeding/Sprigging Chart in the label for this situation.

For Giant Foxtails, Green Foxtail, Yellow Foxtail, Broadleaf Signalgrass Control: Under certain conditions applications of EJECT 75DF herbicide made to annual grasses at 2-4 tiller may not provide complete control. A sequential application will be required for grass control in these situations. Optimum control is achieved when applications of EJECT 75DF herbicide + methylated seed oil is applied either before second tiller or as weed grasses mature.

Biotypes of large and smooth crabgrass in California have shown varied response to EJECT 75DF herbicide. If control failure occurs following the application of a full or split application rates do not reapply EJECT 75DF herbicide, and change to a herbicide with a different mode of action.

For Kikuyugrass, Common Dandelion, English Daisy Control: Tank mix partner or sequential application required.

For Kikuyugrass, Torpedograss Control: Make 3 sequential applications of 0.245 ounce of EJECT 75DF herbicide per 1,000 square feet at 14-21 day intervals.

NOT FOR USE to control English Daisy in California.

Refer in the label for tank mix information and additive rate with methylated seed oil.

Method

[Broadcast](#)

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[Postemergence \(Weed\)](#)