

POME FRUITS (ONLY IN AND AROUND RESIDENTIAL AREA)

General Information

RESTRICTIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Avoid runoff or puddling of irrigation water following application.
- Keep children and pets off treated area until dry.
- Avoid application of ENFORCE 75WSP to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.
- Do not apply more than 8.6 ounces (0.4 lb. of active ingredient) per acre per year.
- Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.
- For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

Limitations, Restrictions, and Exceptions

POME FRUITS

Apply 1.6 ounces (1 packet) per 300 gallons of water

OR

2.1 ounces per acre¹ as foliar spray after petal fall is complete.

- For control of rosy apple aphid, apply prior to leafrolling caused by the pest.
- For first generation leafminer control, make first application as soon as petal fall is complete. Greatest leafminer control will result from the earliest possible application.

For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae.

A second application may be required 10 days later if severe pressure continues, or if generations are overlapping, a single application may result in suppression only.

ENFORCE 75WSP will not control late-stage larvae.

- For San Jose Scale, time applications to the crawler stage. Treat each generation.
- For late-season (preharvest) control of leafhopper species, apply ENFORCE™ 75WSP while most leafhoppers are in the nymphal stage.
- For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.

RESTRICTIONS FOR RESIDENTIAL POME FRUITS:

- Allow 10 or more days between applications.
- Allow at least 7 days between last application and harvest.
- Not for use in California for control on pears.
- Do not apply pre-bloom or during bloom or when bees are foraging.
- The amount of ENFORCE 75WSP required per acre depends on tree size and/or volume of foliage. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees with full foliage.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[After complete petal-fall.](#)

[Rosy Apple Aphid: Prior to leafrolling.](#)