

VEGETABLES: LETTUCE (HEAD AND LEAF)

General Information

- Kills and repels insects, including japanese beetles and over 130 other insect pests.
- Covers up to 16,000 sq. ft. lasts up to 8 weeks.
- Protects trees, shrubs, ornamentals, lawns, structures and listed vegetables from insect damage.
- Just connect to garden hose and spray twist & shoot feature delivers a straight “stream” for taller trees and a “fan” pattern for low lying plants and gardens.

Do not allow children or pets into the treated area until dry.

In New York State, this product may not be applied to lawns within 100 ft. of a coastal marsh, or stream that drains directly into a coastal marsh.

Note: This product is non-staining to most home siding, depending on age and cleanliness. Prior to applying this product to areas that may contact home siding (including vinyl siding), test in a hidden area and recheck in a few hours. If any staining is observed, do not use this product.

Application

Spray as directed - complete coverage is important. Treat when insects begin to appear and repeat treatments to maintain control as directed. Carefully follow all label directions and any limitations of use, such as the amount of spray to apply or PHI (pre-harvest interval) for particular vegetable crops.

TWIST & SHOOT READY TO SPRAY INSTRUCTIONS

- 1) Make sure control knob is in “OFF” position, then connect to garden hose.
- 2) Turn water on at faucet. When spraying low growing plants and small shrubs, twist the control knob right, to the “FAN” position. When spraying taller trees, shrubs and other plants, twist the control knob left to the “STREAM” position for extended reach and more uniform coverage. The product mixes automatically with

the water as you spray.

3) To stop spraying, turn the control knob lever to the "OFF" position. Turn off water at the faucet and disconnect sprayer from garden hose.

Limitations, Restrictions, and Exceptions

VEGETABLES: LETTUCE (HEAD AND LEAF)

Apply thoroughly as a foliar cover spray to both sides of plant leaves and all plant surfaces where insects are present. For best results, apply when insects first appear or damage occurs. Do not exceed pre-harvest interval (number of days to wait between last application and harvest) for treated vegetables.

Note:

- Adult Stages: Japanese beetle; Vegetable weevil.

Method

[Foliar spray](#)

Pre-Harvest Interval

1 days

Timings

[When insects first appear or damage occurs](#)