

RATS

General Information

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. This product is especially suited for use in wet or damp areas, including river banks, gullies, irrigation ditches, railroad tracks, loading areas, under granaries, along fences, garbage dumps and landfills. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not apply bait directly to ground surface or in grass or other ground cover. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

Limitations, Restrictions, and Exceptions

APPLICATION DIRECTIONS

Rats: Place one to four bait packs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

Rats/Mice:

- Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

USE RESTRICTIONS:

- For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. This product is especially suited for use in wet or damp areas, including river banks, gullies, irrigation ditches, railroad tracks, loading areas, under granaries, along fences, garbage dumps and landfills.
- Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not apply bait directly to ground surface or in grass or other ground cover. Do not broadcast bait.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using waterproof gloves collect and dispose of all dead, exposed animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Method

[Bait](#)

Timings

[N.A.](#)