

RATS AND MICE

General Information

DIRECTIONS FOR USE

USE RESTRICTIONS:

This product is for use indoors and in burrows leading indoors only. This product is for control of Norway rats, roof rats and house mice only inside of homes, industrial, commercial, agricultural, and public buildings, and similar man-made structures and in burrows located along the periphery of buildings and likely entry routes into buildings. DITRAC TRACKING POWDER must be placed in locations inaccessible to children, pets, domestic animals, or wildlife. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food.

SELECTION OF TREATMENT AREAS: Determine dry areas where rats and mice will most likely pick up the powder on their feet or fur and ingest it during grooming. Generally, these areas are along walls, by gnawed openings and burrows, in corners and concealed places, in spaces between floors and walls or in locations where rodents or their signs have been observed. Remove goods piled directly on floor and place on skids. Use boxes or other obstacles to force rodents to travel through constricted areas.

Give special attention to the climbing ability of roof rats. For this species, use suitable PVC, sheet metal, cardboard or similar rigid tubing securely attached to rafters or other horizontal surfaces where rats will pass. Employ tubes long enough to prevent spillage of powder through the ends.

Limitations, Restrictions, and Exceptions

APPLICATION DIRECTIONS: Evenly sprinkle 1/2 oz (3 level teaspoons) of DITRAC TRACKING POWDER in patches 12 inches long and 6 inches wide. Apply the powder into the rodent burrow or holes or within walls with a hand bulb or similar duster. Do not use power dusting devices. Sprinkle the powder in patches in such a manner as to expose the rodent to it. Patch size may be (but not limited to) 6" x 12" and

should be adapted to each situation. For rat burrows that are located along the periphery of buildings and that are likely to serve as routes of entry into these structures, place about 5 grams of tracking powder in each burrow with foot pump duster (about 15 pumps). Close burrows with soil, loose leaves, or paper. Repeat if burrows are reopened. Maintain powder in treated areas for at least 20 days.

FOLLOW-UP: Replace contaminated powder immediately. Wearing waterproof gloves, collect and properly dispose of all visible dead animals and leftover powder, as well as any powder that has become accessible to non-target animals. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment.

Method

[Bait](#)

[Sprinkle](#)

Rates

[field_rates 0](#)

•

Timings

[N.A.](#)