

PINK SNOW MOLD - PREVENTATIVE

General Information

HDH Peroxy works best when diluted with water containing low levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. HDH Peroxy will readily mix with clean, neutral water and does not require agitation. Do not combine with any other pesticide or fertilizer, HDH Peroxy is formulated with a minimal amount of surfactant for plants having waxy or hairy surfaces. Additional surfactant may be added, if needed. The use of additional surfactant is acceptable. HDH Peroxy works by surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted.

HDH Peroxy does not produce any visible residue, distinct odor or deleterious effects to plants or to postharvest commodities when used in accordance with label directions. Do not use at higher than recommended dilution rates as leaf burn may result. Do not apply this product through any irrigation system unless directed by the label; refer to Chemigation Directions for Use.

Refer the label of agricultural crops.

Limitations, Restrictions, and Exceptions

FOR TURF APPLICATIONS

Broad spectrum treatment for control of algae, fungi and bacteria on turf.

For use on all turf types such as commercial turf, lawns, athletic fields and golf course fairways, greens and tees.

Use HDH Peroxy to control fungi such as Anthracnose, Brown Spot, Dollar Spot, Copper Spot, Fairy Ring, Pink Snow Mold, Pythium, Phytophthora, Summer Patch, Rhizoctonia, Scum, Take All Patch, Fusarium Blight, Stripe Smut, Leaf Spot, Algae, Slime Molds and their spores.

HDH Peroxy Acts on Contact

For treatment of turf, use on golf course fairways greens and tees to

control/suppress algae, bacterial and fungal diseases and the odors and conditions that these organisms may cause.

Typical treatment rates involve treating approximately 1000 square feet of lawn area with 3 to 10 gallons of diluted solution. Add a surfactant for best results.

Pink Snow Mold:

Preventative - Apply a dilution of 2-6 fl. oz. HDH Peroxy in 3 to 5 gallons of water per 1000 sq. ft. Apply at 7 day intervals.

NOTE: Spray in early fall to reduce the number of dormant spores. Treat throughout winter. May be applied to frozen ground.

1. Optimum treatment time is early morning or late afternoon.
2. For best results, apply immediately after grass has been cut.
3. Applications can be made during wet or rainy weather.
4. Use spray solution the same day it is prepared, do not store and reuse mixed spray solution.
5. HDH Peroxy can be inject through automatic irrigation systems in turf areas. Refer to Chemigation Directions for Use for specific instructions on using this product through irrigation systems.

Method

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

•

Timings

[Early fall to reduce the number of dormant spores.](#)