

TREE CROPS: FOLIAR SPRAY

General Information

General Information

CHEM-SUPRA IRON is a foliar or soil-applied micronutrient and is non-phytotoxic when used as directed. CHEM-SUPRA IRON is completely available and absorbed by the leaf surface or the root system because of the complexing properties. CHEM-SUPRA IRON is recommended on the following crops: Alfalfa, Almonds, Avocados, Apples, Barley, All beans, Broccoli, Cabbage, Cauliflower, Carrots, Celery, Citrus, Corn, Grapes, Lettuce, Milo, Melons, Nectarines, Tea, Rice, Pears, Peaches, Pecans, Peppers, Plums, Prunes, Potatoes, Peanuts, Sorghum, Soybeans, Sugar beets, Sweet corn, Sugar cane, Strawberries, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops.

Limitations, Restrictions, and Exceptions

Foliar Spray: Tree Crops

- Apply 1 gallon CHEM-SUPRA IRON in 500 gallons of dilute spray. If concentrated spray is used, increase the concentration in direct proportion to dilution. If tree crops are sprayed several times a year, the above amount can be split into more numerous applications. Split applications over the year are considered more beneficial than single applications.

CHEM-SUPRA IRON will disperse in water with little agitation. Many pesticides can be added and applied while spraying CHEM-SUPRA IRON. Follow this mixing sequence: 1. Water 2. CHEM-SUPRA IRON 3. Pesticide When foliar spraying CHEM-SUPRA IRON through conventional sprayers, use a minimum of 20 gallons of water per acre. When foliar spraying CHEM-SUPRA IRON with low volume equipment, 5 gallons of water per acre is usually sufficient. If less water is used, slight burning of the foliage may occur. A maximum of 1/2 gallon per acre per application is recommended. Aerial applications should not exceed 1 quart per gallon of water. For best results spray when the crop is in an active growing state, after irrigation or natural rainfall. Spray early in the morning or late afternoon. Mid-day sprays may not be effective because of excessive moisture evaporation. The addition of 1/2% (total solution) of nitrogen solution, ammonium sulfate or L.B. Urea may aid leaf

absorption.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

•

Timings

[N. A.](#)