

## **FOR BROADCAST APPLICATIONS ON ESTABLISHED GRASSES SUCH AS FAIRWAYS AND HOME LAWNS**

### General Information

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

#### PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

#### PRODUCT INFORMATION:

Mecomec 4 Turf Herbicide may be used on ornamental turfgrass including creeping bentgrass (Washington, Toronto, and Seaside) and Kentucky bluegrass (Merion). Mecomec 4 Turf Herbicide is a selective herbicide for the control of surface creeping or prostrate broadleaf weeds. Because Mecomec 4 Turf Herbicide is relatively slow in action, it will not injure bentgrass and other fine turfgrass when used as directed.

#### WHEN TO APPLY:

Make applications on a warm, calm day when weeds are growing vigorously. To obtain the best results, Mecomec 4 Turf Herbicide must be applied when weeds are most susceptible and conditions are such that no damage to grasses will occur. During periods of drought weeds become more resistant to herbicides. Under these same conditions, fine grasses are more susceptible to Mecomec 4 Turf Herbicide, particularly when temperatures exceed 90°F. Mecomec 4 Turf Herbicide is absorbed by the foliage of broadleaf weeds and is translocated throughout the plant.

#### EQUIPMENT:

Wash equipment thoroughly with water and a wetting agent or liquid detergent immediately after use. Spray out, fill with clean water and leave overnight. Spray out again before storing or using for another product. Traces of the product remaining in the sprayer can cause harm to susceptible crops sprayed later.

#### USE PRECAUTIONS:

Mecomec 4 Turf Herbicide is active at low concentrations. Spray drift of this product to susceptible plants may injure beans, tomatoes, peas, cauliflower, cabbage, grapes, sugar beets, tobacco, cotton, sunflowers, boxelder trees, flowers, ornamentals, and fruit crops.

- Do not apply this product through any type of irrigation system.
- Do not cut or mow grass 2 to 3 days before or after treatment so that maximum uptake and translocation can occur.
- Do not spray if rain is likely, otherwise the product may be washed off before leaf absorption
- Do not water grass for at least 24 hours after application.
- Mecomec 4 Turf Herbicide is not recommended for St. Augustinegrass and centipedegrass.
- Do not use Mecomec 4 Turf Herbicide on newly sown lawns or turf for six months after planting.
- Do not use the first cuttings as a mulch for flowers and vegetables.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.

#### WEEDS CONTROLLED

Moderately Susceptible Weeds: Repeat applications at one-half the recommended dosage rate may give better control. Do not make a second application until 30 days after the first treatment.

#### Limitations, Restrictions, and Exceptions

For broadcast applications on established grasses such as fairways and home lawns: Apply 1.25 fluid ounces of product per 1,000 square feet of turfgrass with spray volumes ranging from 0.5 to 4.0 gallons per 1,000 square feet. Or the equivalent dosage rate is 3.33 to 3.5 pints of product per acre with spray volumes ranging from 22 to 174 gallons per acre.

The maximum application rate is 3.5 pints of product per acre per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.0 pints of product per acre per year.

#### Method

##### Broadcast Spray

#### Timings

Make applications on a warm, calm day when weeds are growing vigorously.  
When weeds are most susceptible and conditions are such that no damage to grasses will occur.