

# **FOLIAR SPRAY - PERENNIAL BROADLEAF WEEDS CONTROLLED**

## General Information

## General Information

SA-50 brand BRUSH KILLER is recommended for the control of unwanted woody plants, vines and broadleaf weeds around homes, cabins, buildings, trails, fences, walkways, rights-of-way, and other non-crop areas.

## General Use Precautions

Do not apply this product directly to, or allow to drift to grapes, vegetable crops, flowers, ornamentals or other desirable foliage. Applications should be made when wind is low and there is little chance of spray drift. Do not spray when the wind is blowing toward desirable plants. Do not apply with nozzles that produce a fine-droplet spray. Measuring utensils and spray equipment should not be reused for purposes other than herbicide application.

Do not use on or around edible food or food crops.

Do not graze treated areas.

## Application Directions

### When to Apply

For best results, apply as a full coverage foliar spray when brush and weeds are fully leafed out and actively growing. Wilting or other symptoms will occur in 1-6 weeks. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness.

## Limitations, Restrictions, and Exceptions

## FOLIAR SPRAY - PERENNIAL BROADLEAF WEEDS CONTROLLED

### When to Apply

- For best results, apply as a full coverage foliar spray when brush and weeds are fully leafed out and actively growing. Wilting or other symptoms will occur in 1-6

weeks. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness.

### Foliage Spray

- Using a hose-end or tank sprayer, mix 8-16 tablespoons (4-8 ounces) in enough water to make one gallon mixed spray. For mature or hard-to-control plants (such as ash, blackgum, choke cherry, elm, maple, oak, pine or winged elm), or during drought conditions, use the highest rate of application indicated. Resprouting may occur the year following treatment when lower rates are used. The addition of a surfactant (such as SA-50 brand Surfactant for Herbicides) enhances herbicidal action and improves distribution of the spray on hard-to-wet plant surfaces. Apply at the rate of one gallon mixed spray to approximately 500 square feet of area. Coverage may vary depending on the amount of foliage to be treated. Using coarse spray pattern and low pressure, apply as thorough foliage spray to wet all leaves. See \"Use Precautions.\"

### Stump Treatment

- To kill stumps and prevent sprouting, apply the undiluted product with a paint brush or backpack sprayer to completely cover the freshly cut surface. Do not reuse paint brush. Wrap in several layers of newspaper and discard in trash.

### Vine Control

- When treating vines such as poison oak, poison ivy, etc. that grow on or around desirable plants, do not allow spray to contact desirable vegetation. Cut the vine first and treat as directed for stump treatment.

### Method

[Foliar spray](#)

[Directed](#)

### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

### Timings

[When brush and weeds are fully leafed out and actively growing.](#)