

STONE FRUITS (PLUMS AND PRUNE) - POWDERY MILDEW

General Information

Product Information

Rally 40WSP fungicide is a systemic, protectant and curative fungicide for the control of specific diseases mentioned on the label. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventive spray program. Do not apply this product in greenhouses.

Use Precautions Fungicide Resistance Management

Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with another fungicide mode of action. After two consecutive applications of Rally 40WSP, another myclobutanil product or another DMI, rotate to a product that is effective on the target pathogen and has a mode of action different from Rally 40WSP.

Apply the alternate products within the intervals specified on the label for Rally 40WSP. Do not apply Rally 40WSP at rates below those specified on the label. If tank mixing, use the full label rate of Rally 40WSP with the full label rates of other products effective on the target pest. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

General Use Directions for Tree Fruits

Best control of labeled diseases is achieved when Rally 40WSP is applied on a 7- to 10-day application schedule.

Rally 40WSP is a systemic fungicide and does not redistribute with rainfall after application. Adjust application equipment spray nozzles to apply a uniform spray throughout the entire tree canopy.

Use the following as guidance in determining the amount of Rally 40WSP to be used

per 100 gallons of spray or per acre. Refer to specific tree fruit use directions to determine actual use rates for the control of labeled diseases.

Limitations, Restrictions, and Exceptions

PEACHES

Reapply at 10- to 14-day intervals until terminal growth ceases.

Preharvest Interval: Applications may be made up to the day of harvest.

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Restricted Entry Interval

24 hours